

07/02/99  
Jc658 U.S. PRO

Jc630 U.S. PRO  
07/02/99

<b>UTILITY PATENT APPLICATION TRANSMITTAL</b>  <i>(Only for new nonprovisional applications under 37 CFR 1.53(b))</i>	Attorney Docket No.	B0192/7010
	First Named Inventor or Application Identifier	
	B0192/7010	
	Express Mail Label No.	EL025156278US
	Date of Deposit	July 2, 1999

<b>APPLICATION ELEMENTS</b> <i>See MPEP chapter 600 concerning utility patent application contents</i>	<b>ADDRESS</b> TO: Assistant Commissioner for Patents Box Patent Application Washington, DC 20231
<div>1. <input type="checkbox"/> Fee Transmittal Form <i>(Submit an original, and a duplicate for fee processing)</i></div> <div>2. <input checked="" type="checkbox"/> Specification [Total pages 75] 58 pages specification 1 pages abstract 16 pages claims 90 claims</div> <div>3. <input checked="" type="checkbox"/> Drawing(s) (35 USC 113) [Total sheets 21] <input checked="" type="checkbox"/> Informal <input type="checkbox"/> Formal [Total drawings 19]</div> <div>4. <input type="checkbox"/> Oath or Declaration [Total pages ] a. <input type="checkbox"/> Newly executed (original or copy) b. <input type="checkbox"/> Copy from a prior application (37 CFR 1.63(d)) <i>(for continuation/divisional with Box 17 completed)</i> [Note Box 5 below] i. <input type="checkbox"/> <u>DELETION OF INVENTOR(S)</u> Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b).</div> <div>5. <input type="checkbox"/> Incorporation by Reference <i>(usable if Box 4b is checked)</i> The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.</div> <div>6. <input type="checkbox"/> Microfiche Computer Program (Appendix)</div> <div>7. <input type="checkbox"/> Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary) a. <input type="checkbox"/> Computer Readable Copy b. <input type="checkbox"/> Paper Copy (identical to computer copy) c. <input type="checkbox"/> Statement verifying identity of above copies</div>	
<b>ACCOMPANYING APPLICATION PARTS</b>	
<div>8. <input type="checkbox"/> Assignment Papers (cover sheet &amp; documents(s))</div> <div>9. <input type="checkbox"/> 37 CFR 3.73(b) Statement <input type="checkbox"/> Power of Attorney <i>(when there is an assignee)</i></div> <div>10. <input type="checkbox"/> English Translation of Document <i>(if applicable)</i></div> <div>11. <input type="checkbox"/> Information Disclosure <input type="checkbox"/> Copies of IDS Statement (IDS)/PTO-1449 Citations</div> <div>12. <input type="checkbox"/> Preliminary Amendment</div> <div>13. <input checked="" type="checkbox"/> Return Receipt Postcard (MPEP 503) <i>(Should be specifically itemized)</i></div> <div>14. <input type="checkbox"/> Small Entity <input type="checkbox"/> Statement filed in prior Statement(s) application, Status still proper and desired</div> <div>15. <input type="checkbox"/> Certified Copy of Priority Document(s) <i>(if foreign priority is claimed)</i></div>	
16. Other:	

17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:

- ☐ Continuation    ☐ Divisional    ☐ Continuation-in-part (CIP)    of prior application No.:
- ☐ Cancel in this application original claims of the prior application before calculating the filing fee.
- ☐ Amend the specification by inserting before the first line the sentence:


This application is a ☐ continuation ☐ divisional of application serial no. , filed , entitled , and now .

**18. CORRESPONDENCE ADDRESS**

Correspondence address below

ATTORNEY'S NAME	John R. Van Amsterdam, Reg. No. 40,212				
NAME	Wolf, Greenfield & Sacks, P.C.				
ADDRESS	600 Atlantic Avenue				
CITY	Boston	STATE	MA	ZIP	02210
COUNTRY	USA	TELEPHONE	(617) 720-3500	FAX	(617) 720-2441

**19. SIGNATURE OF APPLICANT, ATTORNEY, OR AGENT REQUIRED**

NAME	Alan W. Steele, Reg. No. P-45,128
SIGNATURE	
DATE	July 2, 1999

CHARACTERISATION OF GENE FUNCTION  
USING DOUBLE STRANDED RNA INHIBITION

The present invention is concerned with  
5 characterization or identification of gene function  
using double stranded RNA inhibition (dsRNAi) and  
methods of identifying DNA responsible for inducing a  
specific phenotype in a cell and a method of assigning  
function to known gene sequences.

10 It has recently been described in Nature Vol 391,  
pp.806-811, February 98, that introducing double  
stranded RNA into a cell results in potent and  
specific interference with expression of endogenous  
genes in the cell and which interference is  
15 substantially more effective than providing either RNA  
strand individually as proposed in antisense  
technology. This specific reduction of the activity  
of the gene was also found to occur in the nematode  
worm *Caenorhabditis elegans* (*C. elegans*) when the RNA  
20 was introduced into the genome or body cavity of the  
worm.

The present inventors have utilized this  
technique and applied it further to devise novel and  
inventive methods of assigning functions to genes or  
25 DNA fragments, which have been sequenced in various  
projects, such as, for example, the human genome  
project and which have yet to be accorded a particular  
function and for use in identifying DNA responsible  
for conferring a particular phenotype.

30 Therefore, according to a first aspect of the  
present invention there is provided a method of  
identifying DNA responsible for conferring a phenotype  
in a cell which method comprises a) constructing a  
cDNA or genomic library of the DNA of said cell in an  
35 orientation relative to a promoter(s) capable of

promoting transcription of said cDNA or DNA to double stranded (ds) RNA upon binding of an appropriate transcription factor to said promoter(s), b) introducing said library into one or more of said  
5 cells comprising said transcription factor, and c) identifying and isolating a desired phenotype of said cell comprising said library and identifying the DNA or cDNA fragment from said library responsible for conferring said phenotype.

10 In a preferred embodiment of the invention the library may be organised into hierarchical pools as described in more detail in the examples provided, prior to step b) such as to include, for example, gene families.

15 According to a further aspect of the invention there is also provided a method of assigning function to a known DNA sequence which method comprises a) identifying a homologue(s) of said DNA in a cell, b) isolating the relevant DNA homologue(s) or a fragment  
20 thereof from said cell, c) cloning said homologue or fragment into an appropriate vector in an orientation relative to a promoter(s) capable of promoting transcription of dsRNA upon binding of an appropriate transcription factor to said promoters, d) introducing  
25 said vector into said cell from step a) comprising said transcription factor, and e) identifying the phenotype of said cell compared to wild type.

In each aspect of the invention, the nucleotide or DNA sequence may either be provided in a sense and  
30 an antisense orientation relative to a single promoter which has the properties defined above, or alternatively it may be provided between two identical promoters. In both embodiments dsRNA is provided from the transcription initiated from the promoter  
35 following binding of its appropriate transcription

factor.

The cell according to the invention may be derived from or contained in an organism. Where the cell is contained within an organism, the organism may be adapted to express the appropriate transcription factor. The organism may be any of a plant, animal, fungus or yeast but preferably may be the nematode worm *C. elegans*, which may be any of a wild type, a *nuc-1* or *pha-ts* mutant of *C. elegans* or a combination of said mutations. In an alternative embodiment the DNA or cDNA library or the DNA homologue or fragment thereof may, advantageously, be transfected or transformed into a microorganism, such as a bacterial or yeast cell, which may be fed to the organism, which is preferably the nematode worm *C. elegans*. In this embodiment of the invention the microorganism may be adapted to express the appropriate transcription factor. Preferably, the microorganism is *E. coli*.

In each aspect of the invention, the DNA library, DNA homologue or DNA fragment may be constructed in a suitable DNA vector which comprises a sequence of nucleotides which encode said transcription factor. Alternatively, said transcription factor is encoded by a further vector. In an even further alternative, the cell or organism may express or be adapted to express said transcription factor. Preferably, any of the vectors used in the method according to the invention comprises a selectable marker which may be, for example, a nucleotide sequence encoding *sup-35* or a fragment thereof. The nucleotide sequence may be orientated relative to a promoter such that binding of a transcription factor to the promoter initiates transcription of the DNA into double stranded RNA. Figure 10 illustrates the vectors and the orientation of the DNA sequence which enable double stranded RNA

production in *C. elegans*. Thus in one embodiment the DNA is located between two promoters on a vector capable of expressing dsRNA upon binding of an appropriate transcription factor to said promoters.

5 Alternatively, the vector comprises two copies of the DNA sequence organised in a sense and antisense orientation relative to the promoter and which marker is selectable when contained in a *pha-1* mutant *C. elegans*. Preferably, the promoters are any of T7, T3  
10 or SP6 promoters and the transcription factor comprises the appropriate polymerase.

Preferably, the selectable marker comprises a nucleotide sequence capable of inhibiting or preventing expression of a gene in said cell and which  
15 gene is responsible for conferring a known phenotype. This nucleotide sequence may be part of or identical to said gene conferring said phenotype, and which nucleotide sequence is itself oriented relative to a suitable promoter(s) capable of initiating  
20 transcription of double stranded RNA upon binding of an appropriate transcription factor to said promoter(s). Alternatively, the nucleotide sequence may be a part of or identical to said gene sequence conferring said phenotype, and which nucleotide  
25 sequence is such as to permit integration of said suitable or further vector by homologous recombination in the genome of said cell and following said integration said nucleotide sequence is capable of inhibiting expression of said gene sequence conferring  
30 said phenotype. In this embodiment said nucleotide sequence comprises stop codons sufficient to prevent translation of said nucleotide sequence following its integration into said genome.

Compounds can, advantageously, in said method be  
35 added to said cell or organism for the purposes of

screening for desired phenotypes, such as for example, resistance or sensitivity to the compound when compared to wild type. The promoters are preferably inducible. The transcription factor may in some  
5       embodiments be phage derived, such as for example, a T7 polymerase driven by a phage promoter. However, when *C. elegans* is utilised a worm specific or tissue specific promoter can be used, such as for example, let858, SERCA, UL6, myo-2 or myo-3. Preferably, the *E.*  
10       *coil* strain is an RNAaseIII and even more preferably an Rnase negative strain.

A further aspect of the present invention provides a method of generating a transgenic non-human organism comprising an exogenous transcription factor  
15       and a transgene comprising a promoter operably linked to DNA fragment which is expressed upon binding of said transcription factor thereto, the method comprising a) providing a first transgenic organism comprising a first construct incorporating DNA  
20       encoding an exogenous transcription factor and a second transgenic organism comprising a second construct including at least one promoter operably linked to a desired DNA sequence which is expressed upon binding of the transcription factor of said first  
25       transgenic organism thereto b) crossing said first and second transgenic organisms and selecting offspring expressing said desired DNA sequence. In one embodiment said first and second transgenic organisms are generated by transforming said first and second  
30       constructs into respective microorganisms for subsequent feeding to the respective organism. Preferably, said second construct comprises said desired DNA sequence in an orientation relative to said promoter so as to be capable of initiating  
35       transcription of said DNA to dsRNA upon binding of

said transcription factor thereto. In this embodiment said second construct comprises two promoters flanking said desired DNA sequence which promoters can initiate transcription of said DNA sequence to dsRNA upon

5 binding of said transcription factor to said promoters. Alternatively, said DNA sequence is provided in a sense and an antisense orientation relative to said promoter so as to produce dsRNA upon binding of the transcription factor to the promoters..

10 In each of these embodiments the first and/or second constructs may preferably be provided with a reporter gene operably linked to a promoter which is capable of initiating transcription of said reporter upon binding of said transcription factor thereto. Preferably, the reporter gene encodes any of Luciferase, Green  
15 Fluorescent protein,  $\beta$  galactosidase or  $\beta$ -lactamase.

The present invention also includes a method of validating clones identified in yeast two hybrid vector experiments which experiments are well known to  
20 those skilled in the art and which experiments were first proposed by Chien et al. (1991) to detect protein - protein interactions. The method according to the invention comprises providing a construct including the DNA encoding a protein identified in a  
25 two hybrid vector experiment, which construct is such that said DNA is provided in an orientation relative to one or more promoters capable of promoting transcription of said DNA to double stranded RNA upon binding of an appropriate transcription factor to said  
30 promoters, transforming a cell, such as a bacterial cell or alternatively transforming an organism comprising said transcription factor with said constructs and identifying a phenotypic change in said cell or organism, which may be *C. elegans* or the like,  
35 compared to wild type. Preferably, the transcription



factor is inducible in the cell or organism. Once again the DNA sequence may be located between two promoters or in both a sense and antisense orientation relative to a single promoter, as described above.

5 Preferably, the promoter is a phage polymerase promoter and said transcription factor is a RNA polymerase, and preferably T7 polymerases. Also encompassed with the scope of the present invention are vectors used to transform said cells or organisms  
10 and the cells or organisms themselves.

In a further aspect of the present invention there is provided a method of alleviating pest infestation of plants, which method comprises a)  
15 identifying a DNA sequence from said pest which is critical either for its survival, growth, proliferation or reproduction, b) cloning said sequence from step a) or a fragment thereof in a suitable vector relative to one or more promoters capable of transcribing said sequence to RNA or dsRNA  
20 upon binding of an appropriate transcription factor to said promoters, and c) introducing said vector into the plant.

Thus, advantageously, the method according to the invention provides a particularly selective mechanism  
25 for alleviating pest infestation, and in some cases parasitic infestation of plants, such that when the pest feeds on the plant it will digest the expressed dsRNA in the plant thus inhibiting the expression of the DNA within the pest which is critical for its  
30 growth, survival, proliferation or reproduction. In a preferred embodiment, the pest may be any of *Tylenchulus* ssp., *Radopholus* ssp., *Rhadinaphelenchus* ssp., *Heterodera* ssp., *Rotylenchulus* ssp., *Pratylenchus* ssp., *Belonolaimus* ssp., *Canjanus* ssp.,  
35 *Meloidogyne* ssp., *Globodera* ssp., *Nacobbus* ssp.,

Ditylenchus ssp., Aphelenchoides ssp., Hirschmenniella ssp., Anguina ssp., Hoplolaimus ssp., Heliotylenchus ssp., Criconemellasp., Xiphinemassp., Longidorus ssp., Trichodorus ssp., Paratrichodorus ssp.,

5 Aphelenchs ssp. The DNA sequence or fragment thereof according to this aspect of the invention may be cloned between two tissue specific promoters, such as two root specific promoters.

10 A further aspect of the invention concerns the vector used in each of the methods of the invention for constructing said library, which vector comprises two identical promoters oriented such that they are capable of initiating transcription of DNA sequence located between said promoters to dsRNA upon binding  
15 of an appropriate transcription factor to said promoters. The DNA sequence may, for example, include a multiple cloning site. Preferably, the expression vector comprises a nucleotide sequence encoding a selectable marker. In one embodiment the nucleotide sequence encoding said selectable marker is located  
20 between two identical promoters oriented such that they are capable of initiating transcription of DNA located between said promoters to double stranded RNA upon binding of an appropriate transcription factor to  
25 said promoters. Preferably, the selectable marker comprises a nucleotide sequence encoding sup-35, for introduction into *C. elegans* having a pha-1 mutation.

Preferably, the transcription factor comprises either a phage polymerase which binds to its  
30 corresponding promoter or a *C. elegans* specific promoter and even more preferably T7 polymerase. Preferably, the vector includes a multiple cloning site between said identical promoters.

35 In a further aspect of the invention there is provided an expression vector for expressing an

appropriate transcription factor for use in a method according to the invention which vector comprises a sequence of nucleotides encoding said transcription factor operably linked to suitable expression control sequences. Preferably, the expression control sequences include promoters which are inducible, constitutive, general or tissue specific promoters, or combinations thereof. Preferably, the transcription factor comprises a phage polymerase, and preferably T7, T3 or SP6, RNA polymerase.

A further aspect of the invention provides a selection system for identifying transformation of a cell or organism with a vector according to the invention which system comprises a vector according to the invention wherein said selectable marker comprises a nucleotide sequence capable of inhibiting or preventing expression of a gene in said cell or organism which gene is responsible for conferring a known phenotype. Preferably said nucleotide sequence corresponds to a part of or is identical to said gene conferring said known phenotype, and which nucleotide sequence is itself located between two identical promoters capable of initiating transcription of double stranded RNA upon binding of an appropriate transcription factor thereto. Alternatively, the nucleotide sequence comprises a nucleotide sequence which is a part of or identical to said gene sequence which confers a known phenotype on said cell or organism, and which is such that following integration of said vector by homologous recombination in the chromosome of said cell or organism said sequence inhibits expression of said gene sequence conferring said known phenotype. Preferably, according to this embodiment the nucleotide sequence comprises stop codons sufficient

to prevent translation of the nucleotide sequence following integration into said chromosome.

Preferably, the known gene sequence comprises a sup-35 gene or a fragment thereof which is selectable by  
5 identifying offspring growing at a temperature above 25°C following introduction in a pha-1 et123ts mutant *C. elegans* worm.

In a further aspect of the invention provides said known gene sequence comprises a sup-35 gene or a  
10 fragment thereof which is selectable by identifying offspring growing at a temperature above 25°C following introduction of said vector in a pha-1 et123ts mutant *C. elegans* worm. An even further aspect comprises a method of assigning function to a  
15 DNA sequence of a multicellular organism which method comprises a) providing i) a construct comprising said DNA fragment cloned between two promoters capable of promoting transcription in said multicellular organism, in a multicellular organism capable of  
20 initiating transcription from said promoter; b) identifying the phenotype of said multicellular organism compared to wild type.

The present invention may be more clearly understood by the following examples which are purely  
25 exemplary with reference to the accompanying figures, wherein:

Figure 1 is a nucleotide sequence of plasmid PGN1 in accordance with the present invention.

30 Figure 2 is a nucleotide sequence of plasmid PGN100 in accordance with the present invention.

Figure 3 is a schematic representation of the vectors used and the transformation regime used in  
35 the methods according to the present

invention.

Figure 4 is an illustration of an expression vector used in accordance with the invention.

5

Figure 5 is a schematic illustration of the T7 RNA polymerase expression vectors used for transforming *C. elegans*.

10

Figure 6 is an illustration of plasmid PGN1.

Figure 7 is a diagrammatic representation of an enhanced vector for dsRNA inhibition encoding sup-35 dsRNA.

15

Figure 8 is an illustration of a vector for integration into the genome of *C. elegans*.

20

Figure 9 is an illustration of the position of a DNA sequence(s) relative to a suitable promoter to initiate expression of dsRNA from the DNA sequence(s).

Figure 10 is a representation of plasmid pGN108.

25

Figure 11 is a representation of plasmid pGN105.

Figure 12 is a representation of plasmid pGN400.

30

Figure 13 is a representation of plasmid pGN401.

Figure 14 is a representation of plasmid pGN110.

35

Figure 15 is a representation of plasmid pAS2 with forward and reverse T7/T3/SP6 promoters.

Figure 16 is a representation of plasmid pGAD424 with forward and reverse T7/T3/SP6 promoters.

Figure 17 is a representation of plasmid pAS2-cyh2-HA+, both T7-final.

Figure 18 is a representation of plasmid pGAD424-without-FULL-ICE-BOTH-T7.

Figure 19 (a) is a representation of plasmid pGN205 and(b) is a representation of plasmid pGN207.

**Example A: Construction of an ordered and hierarchical pooled cDNA library and applications thereof.**

A random ordered and pooled library:

The vector is an *E. coli* vector harboring two T7 promoters, with a multiple cloning site (MCS) in between. The two promoters are orientated towards each other, and towards the MCS. In the presence of T7 RNA polymerase, expressed in *E. coli*, *C. elegans* or any other organism, RNA will be produced, starting from the two T7 promoters. As these are oriented in the opposite sense, both strands of RNA will be produced from the DNA inserted (cloned) into the MCS in between the two promoters which results in the generation of double stranded RNA (dsRNA) upon binding of the T7 RNA polymerase thereto.

A *C. elegans* cDNA library is constructed in the MCS using standard molecular biological techniques. The library is transformed into *E. coli*, and the resulting *E. coli* are grown in culture and stored in 96 multi-well plates. At this stage, plasmid DNA can

be isolated and stored in 96-multi-well plates corresponding to those of the *E. coli* colonies. Approximately 100,000 colonies are scored. In this way, the library will harbor approximately 5 times the total expressed cDNA variation of *C. elegans*, which gives the opportunity for low expressed sequences to be present in the library. This will result in approximately 1041 96-well plates. The plates are hierarchical pooled as necessary. For the present pooling of the clones is arranged in a range of 10 to 100. If the hierarchical pooling is per 8 or 12 (numbers are more convenient as 96-well plates have a 8 to 12 grid), this will result in approximately 87 multi-well plates and approximately 8352 wells. If hierarchical pooling is per 96 wells, which is a full plate, this results in approximately 11 plates and approximately 1041 wells. At any stage of the hierarchical pooling, plasmid DNA can be isolated, which would be less elaborate as less plates are used, but will result in a loss of complexity although this should not be the case in the pooling per 12. The pooling of the DNA can also be carried out with the original DNA.

The experiments below describe how the hierarchical pooling should be performed, both for the DNA and for the *E. coli* library.

An ordered library for RNAi technology, harboring every gene of the *C. elegans* genome, with applications thereof

As the genome-sequencing project is coming to an end, this information can be used in the application of T7 RNA inhibition technology. Every gene of the *C. elegans* genome can be cloned using PCR technology. In

preference, exons will be cloned with a minimal length of 500 bp. If the exons are too small, smaller fragments will be isolated with PCR, or even parts of introns and neighboring exons will be isolated with PCR technology so that at least a sufficient part of the translated region of the gene is cloned. For this, at least 17000 PCR reactions need to be performed. This collection of PCR products will be cloned in a T7 vector as described (two T7 promoters oriented towards each other with a multiple cloning site in between). Every PCR product is cloned independently, or can be used to generate a random library, analogous to the described cDNA library. If every PCR product is cloned individually, the resulting bacteria and plasmid DNA can be pooled in various ways. Firstly, this collection of individually cloned PCR products in the T7 RNAi vector can be pooled randomly, as described in the random library. This pooling can also be done in a more rational way. For instance, the genes of the *C. elegans* genome can be analyzed using bioinformatic tools (in silico biology). Various genes of the genome will belong to a gene family, or will have homologues in the genome. These members of the gene family will be pooled, or the members, being homologues will be pooled. In this way the total number of about 17000 clones is reduced to a more useable quantity. This library can be used to screen for phenotypes in the methods according to the invention. The resulting phenotype gives a functional description to the gene or gene family or gene homologues of the *C. elegans* genome. As the library consists of a part of every gene in the genome, this method enables description of the full genome in functional-phenotypic terms. For this the double



stranded RNA (dsRNA) needs to be introduced in the worm. This introduction of clones alone, or pooled clones, being random pooling or rational pooling can be achieved in several ways as described.

5

**Example of a vector for the expression of double stranded RNAi**

Any vector containing a T7 promoter may be used, and which contains a multiple cloning site (there are many commercially available). Primers containing the T7 promoter and a primer with the reverse complementary strand, both with the appropriate ends are designed. These primers can be hybridized, and if well designed, cloned in the vector of choice. The minimal sequence for a T7 promoter is TAATACGACTCACTATAGGGCGA. Although any vector can be used for the construction of a T7 expression vector there follows an example of how to achieve this with the vector pGEM-3zf(-).

- Vector pGEM-3zf(+) (PROMEGA) was digested with HindIII and SalI
- Primers oGN1 and oGN2 were mixed together at a final concentration of 1  $\mu$ g/30  $\mu$ l boiled and cooled slowly to room temperature.
- The primer was ligated into the vector using standard ligation procedures. The resulting vector is pGN1 (shown in Figure 1) and contains two T7 promoters oriented towards each other, and harbors a multiple cloning site in between.

Sequences of oGN1 and oGN2 are:

- oGN1: AGC TGT AAT ACG ACT CAC TAT AGG GCG AGA AGC TT
- oGN2: TCG AAA GCT TCT CGC ATA ATA GTG AGT CGT ATT AC

35

### Example of the construction of a library

RNA may be isolated from every organism that is sensitive to RNAi. In general the isolated RNA is then  
5 copied into double stranded cDNA, and subsequently prepared in suitable vectors for cloning. Several procedures exist and molecular biology kits can be purchased from various firms including promega, clontech, boehringer Mannheim, BRL, etc which enable:

10

- isolation of RNA,
- eventually polyA RNA can be isolated (several techniques and kits available)
- first strand synthesis with AMV reverse  
15 transcriptase, random hexameric primers and/or oligo (dT) primer
- second strand synthesis with Rnase H, DNA Polymerase I,
- flush ends with T4 DNA Polymerase
- 20 - addition of an adaptor with T4 DNA ligase.
- eventually treatment with T4 polynucleotide Kinase
- cloning of the cDNA into the vector.

The resulting ligation mixture can be considered  
25 as the cDNA library. The ligation contains all cDNA of the procedure ligated into the vector of interest. To order the library, the ligation needs to be transformed into *E. coli* strains.

30

### Application of this *E. coli* or DNA library

T7 RNA producing strain:

- a standard strain is BL21 (DE3): F-ompT[lon]hsds(r-  
m-; and *E. coli* B strain)  $\lambda$  (DE3). Eventually  
35 variants of BL21 (DE3) can be used, although BL21

(DE3)pLysS is used.

- any other *E. coli* strain which produces the T7 RNA polymerase, which may be available needs to be constructed. This can be generated easily using a phage, which is commercially available, in this case, the  $\lambda$ CE6 vector (provided by Promega) is used. Almost every *E. coli* strain can be transfected with this phage and will produce T7 RNA polymerase.

- a RNaseIII mutant *E. coli*:

Various strains are in principle available, we chose in a first experiment to use strain AB301-105: rna-19, suc-11, bio-3, gdhA2, his95, rnc-105, relA1, spoT1, metB1. (Kinder et al. 1973 Mol. Gen. Genet 126:53), but other strains may suit better. This strain is infected with  $\lambda$ CE6 and so a T7 producing variant will be constructed.

Wild type *C. elegans* worms can be grown on the bacteria pools. The bacteria is expressing the T7 RNA polymerase. This results in large quantities of dsRNA in the gut of the *C. elegans*, which will diffuse in the organism and results in the inhibition of expression. This library can now be used for the screening of several phenotypes. This technique has the advantage that it is a much faster to detect relevant genes in certain pathways, than the known *C. elegans* technology. Moreover, if an interesting phenotype is found, the responsible gene can be cloned easily.

Using the hierarchical pooling one can easily find in a second screen the relevant clone of the pool. The inserted DNA of this clone can then be sequenced. This experiment results in genetic and biochemical DATA in one step.

Wild type *C. elegans* strains can be combined with compounds to screen for phenotype, drug resistance and

or drug sensibility. The *C. elegans* strain can be a mutant strain, screening for an enhanced phenotype, reduced phenotype, or a new phenotype. The *C. elegans* strain can be a mutant strain, and the library screen can be combined with compounds. So one can screen for drug resistance, drug sensibility, enhanced phenotype, reduced phenotype, or a new phenotype. The *E. coli* strain may be any T7 RNA polymerase expressing strain, like BL21 (DE3), for example, but the formation of double strand RNA may be enhanced by using a special *E. coli* strain that is RNaseIII negative. RNaseIII recognizes specific loops in dsRNA. Eventually, an *E. coli* strain can be used that is deleted in RNases other than RNaseIII or an *E. coli* can be used that is deleted in one or more RNases. The expression of the T7 RNA polymerase in most known *E. coli* strains and constructs which are available to generate T7 RNA polymerase producing *E. coli* strains, generally comprise an inducible promoter. In this way the production of the T7 RNA polymerase is regulated, and thus the production of the dsRNA. Advantageously, this feature can be used to "pulse" feed the *C. elegans* worms at specific stages of growth. The worms are grown on the non-induced *E. coli* strains. When the worm has reached the stage of interest, the T7 RNA production in the bacteria is induced. This allows the studying of the function of any gene at any point in the life cycle of the animal.

30           Screening the library for homologues of putative  
              interesting human genes, and assign function to  
              these genes

              Hundreds of genes have been isolated in various  
35           projects, being genomic projects, differential

expressed arrays, hybridization studies, etc. The described cDNA library can provide a way to validate and or assign function to these genes in a fast and efficient manner. First of all the worm homologue or homologues or the genes need to be identified by bioinformatic tools (in silico biology). PCR primers are developed and the cDNA fragment is isolated using PCR technology. PCR can be performed on the hierarchical pools. The positive pool or individual wells harboring the bacteria that has the appropriate cDNA is fed to *C. elegans* and the phenotype is scored.

PCR can be performed on cDNA isolated from *C. elegans*. The resulting DNA can be cloned in the T7 vector and transformed in the dsRNA producing *E. coli* on which the *C. elegans* worms are then fed. Depending on which way is faster and more reliable a choice needs to be made.

If the gene belongs to a gene family, the worm may need to be fed on a mixture of bacteria. Each of them harboring a part of the member of the gene family. *E. coli* strains, growth conditions, combinations with compounds can be performed as described above.

If the library rational is used, in which all the genes of *C. elegans* are cloned in a organized and structured way, the *C. elegans* homologue and eventually the other homologues, orthologues, and members of the gene family can be traced back easily in the library using a silico biology. No PCR is involved in this step, and the bacteria and or DNA can be isolated on which the worm will be grown.

### Examples

The idea of the series of experiments was to test

both the RNAi vector and the various *E. coli* strains that were constructed.

1) Construction of a test plasmid

5 Any cDNA that gives a clear phenotype in the worm when knocked-out, or used in a RNAi experiment can be used. It is known that *unc-22* is a good candidate, but many other genes are possible. We opted for a sensitive system that can be used at a later stage.  
10 The system was tested with *sup-35* in a *pha-1* background. Exon 5 of the *sup-35* was isolated by PCR and cloned in the T7 promoter vector pGN1. The resulting vector was designated pGN2. *pha-1* (e2123) mutant worms cannot produce offspring at temperatures  
15 higher than 25°C. This is due to a developmental problem in embryogenesis. When *sup-35* is knocked-out, or inhibited in this strain, offspring may grow at this temperature. Combination of *pha-1* mutant worms and *sup-35* RNAi is a good system to validate the  
20 various options.

2) Testing the RNAi using an *E. coli* strain that produces dsRNA.

- pGN2 was introduced in *E. coli* strain BL21(DE3) and  
25 T7 RNA polymerase was induced with IPTG. *C. elegans* worms (*pha-1* (e2123)) were inoculated on this bacteria, and grown at the restricted temperature of 25°C. As this mutant is an embryonic mutant at this temperature, no offspring will be observed. If the  
30 *sup-35* gene is efficiently inhibited by the dsRNA present in the *E. coli*, offspring will be observed.  
- pGN2 was introduced in *E. coli* strain AB301-105(DE3) and T7 RNA polymerase was induced with IPTG. *C. elegans* worms (*pha-1* (e2123)) were inoculated on this  
35 bacteria, and grown at the restricted temperature of

25°C. As this mutant is an embryonic mutant at this temperature, no offspring will be observed. If the sup-35 gene is efficiently inhibited by the dsRNA present in the *E. coli*, offspring will be observed.

5

3) Improving the worm strain for better uptake of dsRNA.

Before plating the pha-1 *C. elegans* on the *E. coli* strain that produce the double stranded sup-35 RNA. The worm was mutagenised with EMS (Methane sulfonic Acid Ethyl). The offspring of this mutagenised worm is then plated on the bacteria. The worm that feed on this bacteria give larger offspring which has a mutation that results in an improvement of dsRNA uptake, and can be used for further experiments.

10  
15

Stable integration of the dsRNA producing vector into the genome of the T7 RNA polymerase producing worm

20

An *E. coli* vector can be constructed harboring the following features; Two T7 promoters directed towards each other, with a restriction site or a multiple cloning site in between. Furthermore, the vector may contain the *C. elegans* sup35 genomic DNA, engineered in such a way that it contains several stopcodons at various intervals, so that no full length protein can be expressed from the sup35 genomic DNA fragment as illustrated in Figure 8. Any cDNA or cDNA fragment can be cloned in the multiple cloning site between the two T7 promoters. When this vector is introduced in a *C. elegans* strain which expresses T7 RNA polymerase, the cDNA or DNA fragment cloned between the two T7 promoters will be transcribed, generating dsRNA from the cloned fragment.

25  
30  
35

The vector is designed to be used in pha-1 (e2123) mutant worms expressing T7 RNA polymerase. The expression of the T7 RNA polymerase may be constitutive or regulated, general or tissue specific. These pha-1 (e2123) worms cannot produce offspring at temperatures higher than 25°C, which is due to a development problem in embryogenesis. When sup-35 is inhibited or knocked-out in this strain, offspring may grow at this temperature.

When the vector is introduced in the worm, the vector may integrate by homologous recombination (Campbell-like integration). It has been shown that homologous recombination occurs in *C. elegans*, although at low frequencies (Plasterk and Groenen, EMBO J. 11:287-290, 1992). Homologous recombination at the sup35 gene will result in a knock-out of the gene as the two resulting sup-35 genes will harbor the stopcodons. The resulting worm, and its offspring, if this recombination happens in the eggs, will have a copy of the vector integrated in the genome. This can be selected as only the worms for which the sup-35 has been knocked-out will have offspring at temperatures higher than 25°C. Furthermore, the resulting worm will stably produce double stranded RNA from the DNA fragment cloned between the two T7 promoters. This worm can now be considered as a stable transgenic worm strain with a reduction of function of the gene, from which a fragment has been cloned between the two T7 promoters.

The DNA may be delivered to the worm by several techniques, including injection, ballistic transformation, soaking in the DNA solution, feeding with bacteria. New and other methods that increase the transformation efficiencies can be considered.

The target *C. elegans* strain may in addition,



have other mutations than the pha-1 (e2123) mutation,  
and may express other genes than T7 RNA polymerase.

**EXAMPLE B: a Yeast two-hybrid-RNAi vector**

5

A yeast two hybrid vector can be constructed  
harboring the two T7 promoters. The vectors can be  
designed to replicate both in yeast and in *E. coil*.  
In general cDNA libraries for the yeast two hybrid  
10 system are made in the Gal4 or LexA vectors. The  
library is constructed in vectors having the  
activation domain of one of these genes. A vector can  
be constructed that can still perform in the  
yeast two hybrid screen but which also contains two T7  
15 promoters orientated towards each other, with a  
cloning site therein between. The order of the  
sequences in the plasmid will then be "plasmid  
backbone, (GAL4-T7), MCS, T7, backbone". A *C. elegans*  
cDNA library constructed in this vector can be used as  
20 a standard yeast two hybrid library in an experiment  
to isolate interacting proteins with a given protein.  
Once a clone is isolated, the plasmid can be  
introduced in an *E. coil* strain expressing the  
T7 RNA polymerase, and hence will produce dsRNA of the  
25 cloned fragment. The bacteria producing this dsRNA  
can be fed to the worm and phenotypes can be scored.  
As in the previous example, this validation procedure  
for a newly isolated yeast two hybrid clone is  
remarkably shorter than the standard procedure, which  
30 requires PCR and/or cloning steps, RNA experiments  
and/or knock-out experiments. In most cases isolated  
clones are sequenced first, and on the basis of the  
sequence, a decision is made to continue with further  
experiments. In the present invention every isolated  
35 clone can easily be introduced into the appropriate *E.*

*coli* and fed to the worm. Validation is then performed by phenotype analysis.

To apply this procedure a yeast two hybrid was performed using a known gene as bait and the newly constructed library as the target. Proteins coded by the clones in the target that interact with the bait protein, will result in positive yeast clones expressing the reporter molecule such as can be observed by LacZ staining with X-gal. The plasmid coding for the target protein is isolated directly from the yeast strain and introduced in *E. coli*. The *E. coli* is T7 RNA polymerase producing *E. coli*. In this case, double stranded RNA is produced from the DNA cloned in the multiple cloning site of the vector. When this dsRNA is fed to the worm using the methods described previously, the gene has inhibited in the worm, resulting in a particular phenotype.

- This yeast two hybrid vector can advantageously be used to construct an ordered and hierarchically pooled library as described in the previous example.

- A yeast strain can also be constructed that conditionally produces T7 RNA polymerase. After yeast two hybrid experiments, the expression of the T7 polymerase could be induced, resulting in the production of dsRNA in the yeast cell. Consequently the yeast could be fed to the worm. Evidence is available showing that the *C. elegans* worms can feed on yeast.

### Construction of a T7 RNA polymerase producing strain, and applications thereof

A *C. elegans* strain can be constructed that expresses T7 RNA polymerase. The expression can be general and constitutive, but could also be regulated

under a tissue specific promoter, an inducible promoter, or a temporal promoter or a promoter that harbors one of these characteristics or combination of characteristics. DNA can be introduced in this *C.*

5 *elegans* strain. This is done either by injection, by shooting with particles, by electroporation or as aforementioned by feeding. If the DNA is a plasmid as described in the previous examples, i.e. a plasmid harboring a cloned cDNA fragment or a PCR fragment  
10 between two flanking T7 promoters, then dsRNA of this cDNA or PCR fragment is formed in the cell or whole organism resulting in down regulation of the corresponding gene. The introduced DNA can have an efficient transient down regulation. The introduced  
15 DNA can form an extrachromosomal array, which array might result in a more catalytic knock-out or reduction of function phenotype. The plasmid might also integrate into the genome of the organism, resulting in the same catalytic knock out or reduction  
20 of function phenotype, but which is stably transmittable.

- Plasmid DNA harboring a cDNA or a part of a cDNA or an EST or an PCR fragment of *C. elegans* cloned between two T7 promoters as described in Examples A) and B)  
25 can be introduced in the T7 RNA polymerase worm, by standard techniques. Phenotypes can be analysed -DNA from an ordered and pooled library as in Example A) can be introduced in the T7 RNA polymerase worm, by standard techniques (injection, shooting). Phenotypes  
30 can be analysed. With the hierarchical pool, the original clone can be found easily.

- The same procedure can be performed with a mutant worm expressing the T7 RNA polymerase. Screening for enhanced, reduced or new phenotypes.

35 - The procedure can be used to enable screening of

compounds. Screening with either a wild-type strain or a mutant strain for enhanced or new phenotypes.

- The DNA could be introduced in the worm by new methods. One of which is the delivery of DNA by *E.*

5 *coli*. In this case the hierarchical pooled library is fed to the animal. To prevent digestion of the *E. coli* DNA in the gut of the nematode, preferentially a DNase deficient *C. elegans* will be used, such as *nuc-1* (e1392). This procedure would be one of the most  
10 interesting as it would be independent of transformation efficiencies of other techniques, and generally faster and less labourious.

## 2) Putative enhancements of the method.

15 - A vector is designed, so that it harbors the *sup-35* cDNA or a part of this cDNA, cloned in between two T7 promoters. The rest of the vector is as described in Examples A) and B). This vector can be introduced into a *pha-1ts* mutant *C. elegans*. A temperature selection  
20 system exists in this case and only those worms which have taken up the DNA and express the double stranded *sup-35* RNA will survive at restricted temperatures. The hierarchical pooled library can be delivered by any method described above.

25 - The vector can be used to construct a library that is introduced in a T7 RNA polymerase expressing *E. coli*. In this case we have an analogous screening as in part A) with an additional screening for worms where the dsRNA of *sup-35* is active.

30 - The DNA and or dsRNA of *sup-35* could be delivered on a different plasmid. For the feeding, both DNA feeding (Example C) or dsRNA feeding Example A) and B), this means that the two plasmids could be present in one bacterium, or that the worm is fed on a mixture of  
35 bacteria, one of which harbors the *sup-35* construct.

### Example of the construction of a T7 RNA producing *C. elegans*

To produce T7 RNA polymerase in the worm, several  
5 possibilities are possible. The T7 polymerase can be  
expressed under various promoters, being inducible  
promoters, constitutive promoters, general promoters  
and tissue (cell) specific promoters, or combinations  
10 of those. Examples of these promoters are the  
heatshock promoter hsp-16, the gut promoter ges 1, the  
promoter from cet858, but also the promoter of dpy 7  
and the promoter element GATA1. In this example the T7  
RNA polymerase is expressed under the control of the  
hsp-16 promoter that is available in the pPD49.78  
15 vector. The T7 RNA polymerase is isolated as a PCR  
product using the primers of GN3 and GN4.

The resulting PCR product is digested with NheI  
and NcoI, as is the vector in which we want to clone,  
being the Fire vector pPD49.78. The resulting vector  
20 is pGN100 illustrated in Figure 2  
oGN3: CAT GGC AGG  
ATG AAC ACG ATT AAC ATC GC oGN4: ATG GCC CCA TGG TTA  
CGG GAA CGC GAA GTC CG pGN100 is included.

The vector is introduced into the worm using  
standard techniques, such as micro injection, for  
25 example.

The following strains were then constructed:

- Wild-type (pGN100)
- 30 -nuc-1 (e1392) (pGN100)
- pha-1 (e2123) (pGN100)
- pha-1; nuc-1 (pGN100)

All of these strains are able to produce T7 RNA  
35 polymerase when temperature induced or alternatively

by metals such as application of heavy cadmium or mercury. The procedure for temperature induction is to shift the animal to a temperature of 30-33°C for at least one hour, then the animal can be shifted back to standard temperatures (15-25°C).

The wild type strain producing T7 RNA polymerase can be used for the production of any RNA in the worm. More specifically, the plasmids from the described libraries can be introduced in these worms, and phenotypes can be scored.

The nuc-1 mutant worm will be used to introduce DNA via bacteria on which the worm feed. As the nuc-1 worm does not digest the DNA, the plasmid DNA can cross the gut wall. If taken up by the cells that produce the T7 RNA polymerase, dsRNA will be produced thus inhibiting the gene from which the RNA was transcribed.

The pha-1 mutant strain that produced T7 RNA polymerase can be used to enhance the procedures as described above. DNA can be introduced by shooting, micro injection or feeding. More specifically this strain can be used for the vectors that produce dsRNA from sup-35 and from the gene of interest, the latter can be a PCR product, a cDNA, or a library as described.

The pha-1; nuc-1 mutant producing T7 RNA polymerase can be used for the bacterial delivery of the DNA. DNA will preferentially be the plasmid that produce dsRNA from both sup-35 and the gene of interest. The worm strain will preferentially produce the T7 RNA polymerase in the gut. Delivery will preferentially happen by feeding the worm on bacteria harboring the plasmid.

### Application of the RNAi technology in plants

Nematodes are responsible a large part of the damage inflicted on plants and more particularly to plants used in the agricultural industry. The RNAi procedures according to the invention can be applied to plants to prevent these parasitic nematodes from feeding longer. In a first step, a DNA fragment is isolated from the parasitic plant nematode that is critical for the animals survival or growth, or to feed or to proliferate. Any gene from which the expression is essential is suitable for this purpose.

A part of this gene, an exon or cDNA is cloned. This DNA fragment can be cloned under the influence of a tissue specific promoter preferably a root specific promoter even more preferably between two root specific promoters. The DNA of the cloned gene under the control of the root specific promoter can be introduced in the plant of interest, using plant transgenic technology. For every parasitic nematode, a different piece of DNA may be required and likewise for every plant race, a different promoter will be needed.

The root will produce RNA or dsRNA from the introduced piece of DNA when root specific promoter is utilised. As the nematode feeds on the plant, the RNA and/or dsRNA will be consumed or ingested by the nematode. The RNA and/or dsRNA can enter the cells of the nematode and perform its inhibitory action on the target DNA. Depending on the nature of the cloned DNA piece of worm, the nematode will not be able to survive, to eat, proliferate, etc in any case preventing the animal of feeding longer on the plant, and thus protecting the plant.

## Construction of a T7 RNA polymerase producing *C. elegans*

To produce a T7 RNA polymerase or other RNA polymerases in animals, and more particularly in nematodes and most particularly in *C. elegans*, several possibilities can be envisaged. The T7 RNA polymerase can be expressed under various promoters. These promoters may be inducible promoters, constitutive promoters, general promoters, tissue specific promoters, or combinations of those.

### Example 1:

#### Construction of a basic vector for expression of T7 polymerase in *C. elegans*

The T7 polymerase coding sequence was PCR amplified from  $\lambda$  CE6 (Novagen, Madison, USA) using the primers oGN26 (ATGGAATTCTTACGCGAACGCGAAGTCCG) and oGN46 (CTCACCGGTAATGAACACGATTAACATCGC), using standard procedures (PCR, A practical A practical approach, 1993, Ed. J. McPherson, et al, IRL Press). The resulting DNA fragment encoding for the T7 RNA polymerase was digested with AgeI and EcoRI and inserted into the Fire vector pPD97.82 digested with AgeI and EcoRI. The resulting construct encodes for an open reading frame of T7 RNA polymerase in fusion with the SV40 large T antigen nuclear localization signal (NLS) with amino acid sequence MTAPKKKRVVPV. This nuclear localization signal sequence is required to translocate the T7 RNA polymerase from the cytoplasm to the nucleus, where it is able to bind to its specific promoters, designated T7 promoters. Upstream of the coding sequence for the T7 polymerase fusion protein is a minimal promoter (myo-2) preceded by a



multiple cloning site (MCS) in which several *C. elegans* promoters can be inserted. This plasmid (pGN105 shown in Figure 11) is a basic T7 RNA polymerase plasmid which enables the expression of T7polymerase in *C. elegans*. Derivatives of this plasmid wherein promoters are cloned into the multiple cloning site, allow for the inducible, constitutive, general and tissue specific expression of T7 RNA polymerase in *C. elegans*, as expression will be regulated by the promoter cloned in the multiple cloning site.

Although not restricted to these examples, for the following promoters it is known that they induce expression in the following tissues.

let-858 (ubiquitous expression), myo-2 (pharynx expression), myo-3 (body wall muscles), egl-15 (vulval muscles), unc-119 (pan-neuron).

Example 2:

Construction of a vector for expression of T7 RNA polymerase in *C. elegans* muscle tissue.

The T7 RNA polymerase coding sequence was PCR amplified from  $\lambda$  CE6 using the primers oGN43 (GCCACCGGTGCGAGCTCATGAACACGATTAACATCGC) and oGN44 (CACTAGTGGGCCCTTACGCGAACGCGAAGTCCG) digested with AgeI/SpeI and inserted in the pGK13 vector digested with AgeI/SpeI. (This vector contains the strong SERCA promoter which drives expression in the pharynx, the vulval muscle, the tail and the body wall muscle). A nuclear localization signal (NLS) of SV40 large T antigen was inserted in front of the T7 polymerase coding sequence by insertion of two overlapping oligo's oGN45 (CCGGATGACTGCTCCAAAGAAGAAGCGTAAGCT) and oGN46 (CTCACCGGTAATGAACACGATTAACATCGC) into the

SacI/AgeI restriction sites. The resulting construct was called pGN108 as shown in Figure 10. Introduction of this plasmid into *C. elegans* results in the expression of T7 RNA polymerase in the pharynx, vulva muscle, tail and body wall muscles.

To test expression and functionality of T7 RNA polymerase in *C. elegans* under the regulation of the SERCA promoter, pGN108, which encodes the T7RNA polymerase under the control of the SERCA promoter was injected into *C. elegans*. A test vector was co-injected. This test vector encodes for GFP under the control of a T7 promoter (pGN401 in Figure 13). The plasmid pGN401 was constructed by inserting two overlapping oligo's oGN41 (CCCGGGATTAATACGACTCACTATA) and oGN42 (CCGGTATAGTGAGTCGTATTAATCCCGGGAGCT) in the SacI/AgeI opened Fire vector pPD97.82. generating a T7 promoter. Furthermore a selection marker was co-injected to select for transformants (rol6, pRF4). The latter selection vector pRF4 is well known to person skilled in the art. Transgenic F1 could easily be isolated as they display the rol 6 phenotype. These transgenic *C. elegans* all expressed GFP in the pharynx, the vulval muscle, the tail and the body wall muscle. This data show clearly that the T7 RNA polymerase is functionally expressed under the regulation of the SERCA promoter, and that the expressed T7 RNA polymerase binds to the T7 promoter present in pGN401 and initiates transcription of the GFP gene, which is then functionally expressed, resulting in fluorescence in the muscle tissues where SERCA is inducing the expression of the T7 RNA polymerase.

Example 3:

Construction of a vector for ubiquitous  
expression of T7 polymerase in C. elegans

5       The NLS-T7 RNA polymerase fusion gene was  
isolated from pGN108 with XmaI/Bsp1201 and cloned into  
the Fire vector pPD103.05 digested with  
XmaI/Bsp1201. This results in a vector wherein the T7  
RNA polymerase is cloned under the regulation of the  
10       let858 promoter. This specific promoter enables the  
expression of T7 RNA polymerase in all tissues. The  
resulting plasmid was named pGN110 (Figure 14).

Example 4:

15       Construction of a vector for T7 RNA polymerase  
mediated expression of DNA fragments, genes, and  
cDNA's under the control of a T7 promoter.

20       The Fire vector pPD97.82 was digested with SacI /  
AgeI and a T7 promoter sequence was generated by  
insertion of two overlapping oligo's oGN41  
(CCCGGGATTAATACGACTCACTATA) and oGN42  
(CCGGTATAGTGAGTCGTATTAATCCCGGAGCT) into the SacI/AgeI/  
restriction endonuclease sites. This construct (pGN400  
25       Figure 12) contains a GFP open reading frame cloned  
between SacI and EcoRI restriction endonuclease sites  
under the regulation of the T7 promoter. Any gene,  
cDNA, or DNA fragment can be cloned in this vector by  
deleting the GFP gene as a AgeI/ SacI fragment and  
30       cloning the DNA fragment of interest into the vector.  
Preferentially the DNA fragment of interest can be  
obtained by PCR amplification, inserting the SacI/AgeI  
sites in the primers. The resulting DNA fragment after  
PCR amplification is the digested and the GFP gene in  
35       pGN400 is replaced by the amplified DNA fragment.

Every vector that contains a T7 promoter could be used for the purpose of T7 RNA polymerase induced expression in *C. elegans*, such as the commercially available pGEM vectors and the pBluescript vectors. 5 This is clearly shown by the pGN401 vector which expresses GFP under the regulation of the T7 promoter in a transgenic *C. elegans* which expresses T7 RNA polymerase.

10 The use of pGN400 has the advantage that the vector includes a 3'UTR fragment from *unc-54* which enhances the transcription or stability of the RNA.

#### Generation of permanent, tissue specific "pseudo knock-out" RNAi *C. elegans* lines

15 At present, gene knock outs in *C. elegans* are obtained after random, large scale mutagenesis and PCR base sib-selection. This method is bulky, very time consuming and tedious. It has been described that 20 introducing double stranded RNA into a cell results in potent and specific interference of expression of endogenous genes. In *C. elegans* gene expression can be down regulated by injection of RNA into the body cavity of the worm, soaking the worm in a solution 25 containing dsRNA or feeding *E. coli* that express dsRNA corresponding to the gene of interest. *C. elegans* cells have the ability to take in dsRNA from their extracellular environment. It has been reported that mRNA is the target of this ds RNA mediated genetic 30 interference (Montgomery and Fire 1998). It is also suggested that the targeted RNA is degraded in the nucleus before translation can occur. Although the RNAi mediated reduction of gene expression can be passed on to the next generations, heritability is 35 poor and the effect is rapidly lost during further

offspring. This is probably due to a continued decrease of the dsRNA pool. We propose here a method to construct *C. elegans* lines with a permanent, inheritable, RNAi phenotype. The method encompasses the generation of transgenic *C. elegans* lines by introducing plasmids containing cDNA fragments of the target gene in the sense and antisense orientation under control of a worm promoter or by transcription of an inverted repeat of the cDNA from a single construct. Alternatively, ds RNA can be transcribed from a vector harboring a cDNA flanked by two T7 promoters in a *C. elegans* strain that expresses T7 polymerase. The result is a transgenic worm with an heritable stable "pseudo knock-out" phenotype. The expression of the cDNA or the T7 polymerase can be general and constitutive but could also be regulated under a tissue specific promoter. In contrast to RNAi induced by external ds RNAi (injected, soaked or feeded) this method would enable to obtain conditional, tissue specific inhibition of gene expression.

Inhibition of unc-22 expression by RNA interference results in a "twitching" phenotype.

25

Unc 22 cDNA (exon 22) was cloned in sense and antisense orientation in pPD103.05. (A. Fire nr L2865) containing the let 858 promoter that is capable of expressing RNA sequences in all tissues. The resulting plasmids were named pGN205 (Figure 19a) and pGN207 (Figure 19 b). These constructs were introduced into *C. elegans* together with a selectable marker (*rol-6*; GFP). Transgenic F1 individuals (expressing *rol-6* or GFP) showed a "twitching" phenotype indicating that RNAi could be mediated by endogenous transcription of

35

RNA from transgenic DNA. The RNAi phenotype co-segregated with the selectable marker into further offspring. This resulted in the generation of C. elegans lines with permanent RNAi phenotype.

5

Generation of stable lines T7 RNA polymerase lines and generation of dual transgenic worms.

10 An expression system in C. elegans based on an exogenous RNA polymerase demands two plasmids. One is encoded for the RNA polymerase under the control of a specific promoter, while the other plasmid encodes for the DNA fragment to be expressed, under the regulation of the T7 promoter. In the case of semi stable RNAi  
15 also designated pseudo stable knockouts, the DNA of interest is cloned between two T7 promoters so that dsRNA can be produced.

As the T7 RNA polymerase expression system is known to be a high expression system this will result  
20 in problems to generate dually transgenic animals. If the gene to be expressed in the C. elegans nematode is toxic, this will result in lethal effects and hence in the construction of a C. elegans without highly regulated stable expression of the gene of interest.  
25 If the gene of interest is essential for the survival of the organism, RNAi with a DNA fragment from this gene will also result in lethal effects, so that pseudo- stable knockouts are not possible.

To overcome this problem the present inventors  
30 have designed a system consisting of two transgenic animals. The first animal is transgenic for the T7 RNA polymerase, This T7 RNA polymerase can be expressed in all cells or specific cells or tissues as has been shown in previous examples. The second transgenic  
35 animal is transgenic for the DNA fragment of interest.

This can be a gene or cDNA linked to a T7 promoter, or if one wants to perform RNAi a DNA fragment of such gene cloned between two T7 promoters.

Both transgenic animals are viable and do not  
5 show any aberrant phenotypes. This is because the T7RNA polymerase expressed in the first transgenic organism is not toxic for the organism, even if expressed at relative high levels. In the second transgenic organism, the gene of interest is not  
10 expressed or the dsRNA is not produced as these transgenic animals do not contain the T7 RNA polymerase.

Expression of the gene or cDNA of interest or RNAi with a DNA fragment can now be obtained by mating  
15 the two transgenic animals. The offspring of these are dually transgenic and express the gene of interest or express dsRNA of the DNA fragment of interest. To generate sufficient males in such a mating, one of the transgenic animals males can be a *C. elegans*  
20 mutant with a phenotype favouring generation of males. An Example of such a mutant is him-5. Preferentially such a mutant will be used to make a *C. elegans* transgenic for T7 RNA polymerase, while the hermaphrodite harbors the DNA fragment under the  
25 regulation of the T7 promoter.

To select efficiently for the dual transgenic offspring a second transgene can be introduced in the second transgenic animal. This transgene contains a reporter gene under the regulation of the T7 promoter.  
30 The reporter gene can be GFP, luciferase, Beta galactosidase, or beta-lactamase. an example of such a ttransgene are the vectors pGN400 and pGN401.

To obtain inducible, tissue specific expression of a transgene in *C. elegans* we can make male stock  
35 (i.e. him-5) carrying the T7 polymerase construct

under the control of different *C. elegans* promoters that enable tissue specific expression such as). This males can be crossed with hermaphrodites carrying the gene of interest under the control of a T7 promoter.

5        Furthermore, the transgenes can be integrated into the genome of the animal. Methods to generate stable integration of a plasmid into the genome of the animal have been described (Methods in cell biology, Vol. 48, 1995, ed. by epstein and Shakes, academic  
10       press) and involve radiation of the animal. This can be done for both animals animals, but preferentially, the animals expressing the T7 RNA polymerase are subject to such traetment. This result in a collection of *C. elegans* nematodes that stably  
15       express T7 RNA polymerase under the control of various promoters. examples of such promoters are the myo-2 (pharynx expression), myo-3 (body wall muscles), egl-15 (vulval muscles), unc-119 (pan-neuron), SERCA (muscles), let858 (all cells) ges-1 (gut).

20

#### Construction of RNAi T7 promoter yeast two hybrid vectors

##### pGAD424 with forward and reverse T7/T3 and or Sp6

25        In most two-hybrid experiments a cDNA library is cloned in plasmid pGAD424 (Figure 16) which has been engineered with additional restriction sites in the polylinker such as a NcoI site (Clontech). This library allows for screening of binding proteins in a  
30        yeast two hybrid experiment. We constructed a new yeast two hybrid vector with the same possibilities to perform yeast two hybrid, but which contain two additional T7 promoters, so that the vector can be used for T7 RNA polymerase induced pseudo-stable  
35        knock-outs. For this we inserted a forward T7 by using



a T7- linker (consisting of the following primers  
aattcttaatacgaactcactatagggcc and  
catgggccctatagtgagtcgtattaag) into the EcoRI-NcoI site  
of pGAD424. The resulting vector was designated  
5 pGAD424-without-FULL-ICE-both-T7. Care was taken to  
eliminate stop codons and using maximal polylinker  
compatible amino acids. We adopted the same strategy  
for the reverse T7 (consisting of both primers  
gatccgtcgacagatctccctatagtgagtcgtattactgca and  
10 gtaatacgaactcactatagggagatctgtcgacg) with BamHI and  
PstI. To avoid loss of SalI, we included this site in  
the primer.

The SalI site is important as most libraries are  
cloned in this site, adapters are available. This  
15 makes the newly constructed vector compatible with  
existing vectors.

pAS2 with forward and reverse T7/T3 and or  
Sp6

20

An analogous yeast two hybrid vector was  
constructed based on pAS2 (Clontech). By partial EcoRV  
digestion we were able to remove a significant part of  
the cyh2 gene. The right construct can be isolated and  
25 checked by a restriction digest with BglII. this  
restriction site is present in the EcoRV fragment of  
PAS2 to be eliminated. This eliminates the cyh2 gene  
which is slightly toxic gene and involved in growth  
retardation. This gene is non-essential for the  
30 performing of RNAi and Yeast two hybrid experiments.  
After the elimination of the EcoRV fragment, The  
EcoRI restriction site which is located between the  
DNA sequence encoding for GAL4DB and HA (epitope)  
becomes unique for the plasmid, and can be used to  
35 substitute HA with a T7 promoter containing linker.

This ensures persistence of all restriction sites, allowing both in frame cloning and compatibility with previous vectors and pGAD424. We used the following linker (primers: aattcttaatacgcactcactatagggca and  
5 tatgccctatagtgcgtcggtattaag) using EcorI and NdeI cloning sites. We adopted the same strategy for the reverse T7 (primers:  
gatccgtcgacagatctccctatagtgcgtcggtattactgca  
catggggccctatagtgcgtcggtattaag and  
10 gtaatacgcactcactatagggagatctgctgcgcg) with BamHI and PstI. To avoid loss of SalI we included it in the primer. The resulting vector was designated pAS2-cyh2-HA+both T7-final.

Having the T7 promoter (or alternatively the T3,  
15 or SP6 promoter) in pGAD424 allows to go quickly from interacting protein to RNAi and assigning function to the isolated DNA fragment. An additional advantage is the ability to make by in vitro transcription coupled to in vitro translation (There is an ATG in frame with  
20 either GAL4DB or GAL4AD) labeled protein which can be used for in vitro controls (e.g. pull down assays) of the actual protein-protein interaction.

The sequences of the plasmids produced and the SP6 and T3 polymerase are identified in the Sequence  
25 Listing provided below:

SP6 DNA-dependent RNA polymerase:

SEQUENCE ID NO. 1

swissprot accession number P06221

protein sequence:

```
1 mqdlhaiqlq leeemfnggi rrfeadqqrq iaagsesdta wnrrllseli apmaegiqay
61 keeyegkkgr apralafllq venevaayit mkvmdmlnt datlqaiams vaeriedqvr
121 fsklegghaak yfekvkkslk asrtksyrha hnvavvaeks vaekdadfdr weawpketql
181 qigtllleil egsvfyngep vfmramrtyg gktiyyllqts esvgqwisaf kehvaqlspa
241 yapcvipprp wrtpfnggfh tekvasrirl vkgnrehvrk ltqkqmpkvy kainalqntq
301 wqinkdvlav ieevirldlg ygvpsfkpli dkenkpanpv pvefqhlrgr elkemlspeq
361 wqqfinwkge carlytaetk rgksaavvr mvqgarkysa fesiyfvyam dsrsrvyvqs
421 stlspqsndl gkallrftg rpvngvealk wfcinganlw gwdkktfdvr vsnvldeefq
481 dmcrdiaadp ltftqwakad apyeflawcf eyaqyldldv egradevrth lpvhqdgscs
541 gighysamlr devgakavnl kpsdapqdiy gavaqvikk nalymdadda ttftsgsvtl
601 sgtelramas awdsigitrs ltkkpvmtp ygstrltcre svidyivdle ekeaqkavae
661 grtankvhpf eddrqdytlp gaaynymtal iwpsisevvk apivamkmir qlarfaakrn
721 eglmytlptg fileqkimat emlrvrctcm gdikmslqve tdivdeaamm gaaapnfvhg
781 hdashliltv celvdkgvts iavihdsfgt hadntltlrk alkqgmvamv idgnalqkll
841 eehevrmvmd tgievpeqge fdlneimdse yvfa
```

T3 DNA dependent RNA polymerase:

SEQUENCE ID NO. 2

swissprot accession number P07659

protein sequence:

```
1 mniieniekn dfseielaai pfntladhyg salakeqlal ehesyelger rflkmlerqa
61 kageiadnaa akpllatllp kltrrivewl eeyaskkgrk psayaplql kpeasafitl
121 kvilasltst nmittiqaag mlgkaiedea rfgrirdlea khfkkhveeq lnkrhgqvyk
181 kafmqvvead migrqllgge awsswdkett mhvgirliem liestglvel qrhnnagnags
241 dhealqlaqe yvdvlakrag alagispmfq pcvvppkpww aitgggywan grrplalvrt
301 hskkglmrye dvympvyka vnlaqntawk inkkvlavvn eivnwknpcv adipslerge
361 lppkpddidt neaalkewkk aaagiyrldk arvsrrisle fmleqankfa skkaiwfpyn
421 mdwrgrvyav pmfnpqgndm tkglitlakg kpigeegfyw lkihganacag vdkvpfperi
481 afiekhvddi lacakdpinn twaeqdsfp cflafcfeya gvthhglsyn cslplafdgs
541 csgiqhfsam lrdevggrav nllpsetvqd iygivaqkvn eilkqdaing tpnemitvtd
601 kdtgeisekl klgtstlaqq wlaygvtrsv tkrsvmtlay gskefgfrqq vlddtiqpai
661 dsgkglmftq pnqaagymak liwdavsvtv vaaveamnw ksaakllae vkdkktkeil
721 rhrcavhwtt pdgfpvwqey rkplqkrldm iflgqfrlqp tintlkdsqi dahkgesgia
781 pnfvhsgdgs hlrmtvvyah ekygiesfal ihdsfgtipa dagklfkavr etmvityenn
841 dvladfysqf adqlhetqld kmpplpkkgn lnldilksc fafa
```

pGN108:

[illegible][illegible]

ttctctcttgactctctctctccgccaataagcctagcattttatgataatmtgattacacacactcagagtctcttcgacatgataaagtgmtcattggcac  
tcgcccttaacagtagcatgacaaaggcggtattatcagtcgatattgaaagacaaactcctcaaatgtgtgctcatttggagcccgctgtggggcagctg  
ctctcaatatattactaggcgagacgaggagggggacctatcgaactcgcgatgagccatttctctttatgactctcttcaactctcacacattaat  
cgattcatagactcccatattcttgatgaagggtgtgggttttttagctctttttccgatttgtaaaagggaaggcgtgacgatgttaggaaaaagagaacg  
gagccgaaaaaacatccgtagtaagtcttcttttaagccgacacttttagacagcattcgcgcgtatggtgaagtttaaaatataaaaattag  
ttcaatttttaattactaataatggcaaaagtttttcaagaactctagaaaaactagcttaattcatgggtactagaaaaattctgttttaaattaattta  
tcttaagatgttaattacgagaagctttttgaaaaatctcaatataaagaattgcccgtatttagaataaaagtcttcagaaaatgagtataaagctcaaataga  
agttgttttttaaaggaaaaacacgaaaaaagaaacactattatcttttctcccgctgaaaaattagttgtgtgataatgtagacgcgtctgtattgtcact  
cgtctcttcacacagctgtctctctcacttagacccaacgaaaaaaaacatcagcttgagacgtgtgaattgctcttcacacagagcctgtctcttcca  
ccagtagacaaaaaaaattgtgctttctttattatgttaggtctacaaaaaaaagtagatcgagttgtgtgtgcgtgtgtctgtctcacaccacctccgc  
ctccgcagcagcacacaatcattctgtgtgtctgcagcttccctgtatgcgcgggtcgaatgcaccactctgacgcgcgaactcacaccacacactc  
actttcgggtgtattactacagctatcgtgtgtctgtctcccgctctttcgtccccactcactctctatttcccctgtgtgtattgttttttaaaatggta  
caccactctgacgtttctactcttctgttttccgtccatttagatttatctggaatttttttaaaatttaggccagagagttctatgtctgtcttaaaagtcta  
ggctcagacatacatttctatttctatcaaaaaaaaagttagataaaagaaaactgtttatcagaaaagagtgtgtctcgttgaaatgattcaaaaaaaatt  
cccacccctcgtgtttctcctcaaaatagatgatcaacggatttttctctctgattcaatttttctgtcgcctctgtctgcctcaaaagtgtgtgtcctcgagcaaa  
agatgagagagaattacaacagaaaatgaaaaaaggttggccaaataatgaagtattacggagatgagggaagatattaatgttctttacgggtttgga  
ggggagagagagatagatttgcattcaaaactcgcctttatcatgtcttttagaaatctaaaatagatttttctatattttaataagaatacagagaata  
cagtaatttgcgaattttctgccaaaaatacacagaaatttgggtgtctgccacgatctcggcttagtgggtcatttgggttttaaaagttaataaaatttcaaa  
ttctagtgtttaattccgcataatggacctaataatgggtttgtctatcttcaacaagaatacgtgaaaatcgtgtgttctgcaatttcttttcaaaaaata  
cacgaaaataatgtaatttccgaaatagagggtctcggccagtttcagtcacagtgccaggagatttatcacgaaaaaagtgtgacctgacatcacaat  
ttccggaataacgcgaatcaatttagttttgtctatctttgaacaaaaaatcagacatccctatctgatttgcgaatttctgtcgtttctctccaaaatga  
cagtcttagaatttaaaatctcgtgcgaactgggacgttagatcttttctcccggttttattttatttattacactgattgactaaagctccacacacag  
ccagttgtgtccatactacacacacacacacacaaatgtcagaaatttatgtgtttatccctgttgaattcgttccgtgtgtctctctctctatctattt  
agccgagaaagctccagagaatggagcacacaggaatccgggcgcgcgatgtctgcgggagatggcgccgcctgggaagccgcggagagatatca  
gggaagatcgtctgatttctctcgatgccacctcactctcaggtttctccgcctgtttactccctgcggaacctgattttccc

SEQUENCE ID NO. 4

aaagcttgcacgctcaggccttgctgactctagacacatttcagctacatgatacatggaatccccgcctcccaatcaccaccacagggaataa  
gaagggtctgcgcgaataaatacaagttatctccaggctcgcgcacccaccaggcgggtgacttctccaccatttcaatcaccctcggggtacg  
ggattggccaaaggacccaazggtatgtttcgaatgatacacaatacatagaacatttcaggaggacccttgcctggagggtacgcagctcagaa  
aaatgactctccaaaggaaagcgtgaagggtaccggtaatgaacacgattaacatcgtctaaagacgacttctgacatcgaactgctgctatccc  
gttcaacactctggctgacattacggtgagcgtttagctcgcgaacagttggcccttgagcatgagcttcatcagatgggtgaagcacgcttcgcaa  
gattgttgagcgtcaacttaaaagctgtgaggttgctgggataacgtctgcgcgaagccttcatcactacccctactccctaagatgattgcacgcatca  
cgactgttgtagcgaagtgaaagctgaagcgcgcgaagcgcgcacacgttccacgtcctgcgaataaactgaagccggaagccgtgacgtcatca  
cctaataagcacactctgcttgccttaacacagctgtgacaatacaacgtctgaggtcgtgtagcaagcgaatcgctggcgccattgagagcaggctgc  
cttcggctgctacgtgaccttgaagctaaagcactcaagaaaacgttggaacaacatcaaacgctgtaggtgcacgtctacaagaaagcattta  
tgcaagttgctgaggctgacatgctctctaaagggtctactcgttgctgcgcgagcgtgcttctgctgataaggaaagactctattcatgtaggagtacgc  
gcatcgagatgctcatgtagtcaaccggaaatggttagcttacacgcgcgaataatgctggcgttagtggtcaagactctgagactatcgaactcgcacc  
gaatacgtgtaggctatcgaaccctgtcaggtgctgctggtcgtgcatctctccgatgttccaaccttgcgtgagttctctaaagccgtggactggcatt  
actggtgtggtgatttggggttaacggtctcgtcctctggcgtggtgctgactcacaagtaagaaagcactgagtcgctacgaagacgtttacatgctt  
gagggtgtacaagcgattaaactgcgcaaacaccgcgatggaatacaagaagagctcgtgctgccaacgttaatacacaagtggaagcat  
tgtccggctggagacatccctgcgattgagcgtgaagaactcccgatgaacccgggaagacatgcacatgaatccctgaggctctcaccgcgtggaaa  
cgtgctcgcgtcgtctgttaccgaacgacaggctcgtcgaagtctgcgttatcgaatgcagctgagttcgtgtgacagccaataagtttgcataccat  
aaggcccatctgcttcttacaacatgtagctgctgcggtctgtttacgcgtgttcaacgttcaacccgcaaggtaacgattgacccaaggaactgctt  
acgtggtgcgaaggtaaaccaatcgtgaaggaaggttactctgctgctgaaatccacggtgcgaactgtcgggtgtcgataagggttcccttcg  
agcgcatacaagttcattgaggaaaaccacgagaacatcatgcttgcgctlaagctccactggagaacacttgggtgggtgtagcaagattctccgtt  
gcttctctgcttctgcttgagtacgttgggtgacagcaccacggcctgagctataactgtctcccttccgctggcgttgacgggtcttctcgtgcatc  
cagcacttcccgatgctcgcgagatgaggtagggtgctgcgcggttaacttgccttagtgagaccgttcaggacatctacgggattgtgctaag  
aaagtcaacgagatttacaagcagacgcaatcaatgggagaccgataacgaagttagttaccgtgaccgatgagaacactgtgtaaatctctgagaaa  
gtcaagctgggcactaaggcactgggtgtgtaagctgctcaggtgttactcgcagttgtactaagcgttcatgctacgctggttctacgggtc  
caaaagatnccgcttccgtcaacaagtgtcgtgaagataccattcagccagctattgattccggcaagggttccgatgttcatcagccgaatcaggctg  
ctgatacatgctgaagtgtgatttgggaatctgtgtagcgtgacggtgtgctgctggttgaagcaatgaactgtgcttaagctgtctgaagctgctg  
ctgtctgaggtcaagagataaagaagactggagagatttccgcaagcgttgcctgtgctgacttggtaacctgatgttccctctgttggcgaggaatac  
agaagccatttcagacgcgttggaacctgtgcttccctgcgttaccgcttaccgcttaccacccaacaaagatacgagattgtagcacaca  
aacaggagcttggatcgtcttcaacttgtacacagccaagacggtagccacctctgaagactgtagtgtgggcacacgagaaagtcaggaaatcga  
atcttgcactgattcagactccttccgttaccattccggctgacgcgtgcgaacctgttcaagacagtgctgcgaaactatgttgacacatatgagtct  
tgtagtactctggtgatttctacgaccagttcgttgaccagttgcacgagcttcaattgacaaaaatgccagcacttccgctaaaggtaactfgaacct  
ccgtgacatcttagatgctggacttccgttccgctgaagaatnccaactgagcgcggctcgttaccattaccaacttctgtgtgctcaaaaaataatagg  
gcccgtgtcatcagagtaagttaaacrtgagtcttactaactaacgagtaataatttaatttcagcatctcgcgcgcgtcctctgacttctaaagtccaatt  
actctcaacactccctacatgctcttctccctgtgctccaccctctattttgttattatcaaaaaactcttcttaatttcttgttttagcttctttaaagta  
cctctcaacaatgaattgtatgatacaaaaaatagaattattctgtaaaaaaagtcgaaaaaaatgtgtctccctcccccattaataataattctatccca  
aaatctacacaagtctgtgacttcttctgttttttcttctgataaatttttttgaaacatcatagaaaaaacccgacacaaaatactctcatatgctt  
acgttcagtttatgacgcgcaatttttcttgcacgtcttgcgcctctcatgctcaacatcgtctcatcgtgaaaaaggttttgagatttttgcgaattt  
caatcaagtgaagttatgaattaaatttccgtcttctgttgggttccctatttcttgaagagtttcgagagacgcttcttctgtctaaatca  
caaglatgtatgagcacgatgcaagaagatcgggaagaggtttgggttgagggtcagtggaaggtgagtagaagttgaattgaaggtggagta  
gtgtctatgggttgggttgccttaaatgacagaatacattcccaataatacaaacataactgttcttactagtcggccgtacgggccccttctcgcgcg  
ttcgggtgatgacggtgaaaaccttgacacatcgagctcccggaagacggtcacagctgtctgtgaagcggatgcccggagcagacaagccgtca  
ggcgcgctcagcgggtgttgccgggtgtcgggggtgctgcttaactatgcggcatcagagcagangtactgagagtgaccataatgcgggtgaaata  
ccgcacagatgctgaaggagaaaaatccgcatcaggcggccttaaggcgcctctgtatagcctatmttataggttaatgtcatgataaataatggttctt  
agacgtcagggtgacatmttcggggaaatgtgcgcggaacccctattgttatttttctaaatacatcaaatatgtatccgctcatgagacataaacct  
gataaatgtcttaataatantgaaaaaggagatgagatgagatncaacatmccgtgtcgccttattcccttmttgcgacatttgccttctgttmttgc  
ccgaaaaacgctggtgaaagtgaagaagatcgtgaagatcgttggttcacagatgggtttacatcgaactgtgactcaacagcggtaagatcgtgag  
agtttgcggccggaagacgtttccaatgtagtagcacatmtaagtttctgtatgttggcgcttattccgttattgacgcggcgcaagacaaactg  
gtcggcgatcacactattctcagaatgacttggtagtactaccagctcacagaaaagcatctacggatggcatgacagtaagagaattatgcagt  
ctgccataacatgagtataacactcgcggccaacttacttctgacaacgacggaggaccgaaggagtaaccgctttttgcacaacatgggggat  
catgtaactcgccttgatcgttgggaaccggagctgaatgaagccataaccaacgacgagctgacaccacgatgctgtagcaatggcaacaacg  
ttgcgcaactatttaactgcgaactacttactgtatcttccggcaacaatttaatagactgtagtgaggcgggataaagtgcaggaccacttctgcgt  
cggcccttccggctggtgttattgtctgataaacttgagccgggtgagcgttgggtctcgcgttactgtcagcacttggggccagatggtgaagccc  
tccgtctgtgtagttatcagacgacggggagtcaggaactatggatgaacgaataagacagatcgtgagatagggtgctcactgattaaagcattg  
gtaactgtcagacgaagttactatcatatcatmttagatgtatmtaaaacttmttaaaaaaggaatcaggtgtagaagatccctttttgtcgtctatcgt  
gaaaaacccataactgttttcttgcacttagcgctcagcttccggatccgaagaagactcaacatcttmttcgaagcttaactgtctcagcagcgcgtga  
taccataatctgtctttagttagcgttagttaggccaccacttcaagaactctgtgacccgcttataactcgtctgtctaatctgttaccagtgt  
gtgtgtccaggtggcgataagtcgtcttaccgggttggaactcaagacgafagtaccggaataaggcgcggctgggtgtaacgggggttgc

[illegible]

pGN400:

SEQUENCE ID NO. 5

aagcttgcacgctgcagggcctggctgactctagacacmttcagctacctagatcatgatacccccgcctcccaatccaccaccagggaaaa  
gaagggctcgcgcaaaaatcaagttatctccagggctcgcgcatcccacgagcgggtgacttctccaccactttcatmaaccctcggggtaag  
ggattggccaaaggacccaaggtargtttgaatgatactaacatacagaaacatmttcagggagggacccctgcttgagggtaccgagctcccg  
gattaatcagactactatccggtagaaaaatgagtaaaaggagaagaaacttttactggagttgtcccaattctgtgaattagatgtgtatgaatg  
ggcacaattttctgtcagtgaggagggtgaggtgatgcaacatacggaaaaacttacccttaattttttgactactgtgaaaaactaccgtgtccatgg  
gtaagtttaaacataatatactactaacctgatttmaaaatttcagccaactgttactacttctgttatgtgtttcaatgcttctcgagataccag  
atcatatgaaacggcagactmttcaagagtgccatgcccgaaggttatgtacaggaaagaaactatattttcaagatgacgggaactacaagacac  
gtaagtttaaacagttcgggtactactaacatacatatttmaaaatttcaggtgtgaggtcaagtttgaaggtgataccctgtttaaagaatcagttaaa  
aggattgtattttaaagaagatggaacattctgtgacacaaattggacataactataactacacaaatgtatatacatatggcagacaaacaaaagaat  
ggaaatcaaggtgtgaattttaaactatgtatttactactaactatctgatttmaaaatttcagaaacttcaaaaattgacacaaactgaagatggaagcgttca  
actagcagaccattatcaacaaaatactccaattggcgatggccctgtcttaccagacaaccattaccgttccacacaaatctgccccttgcgaagatc  
ccaacgaaaaagagagaccacatggtcttctgtgattgttaacagctgctgggattacacatggcatggatgaactatacaaatagcattctgtagaattc  
caactgagcgcggctgctaccattaccaactgtctgtgtgtaaaaaataatggggccgctgtcatcagagtaagtttmaactgagttactactactaa  
cgagtaattttttaaattttcagcattctcgcggcggctgctgtacttcaagtcacacttcaacatccctacatgcttcttccctgtgtctccccc  
cctattttgttatttcaaaaaaacttctttaaattctgttttttgaacttcaacttcaacaaatgaaattgtgtatgtaaaaaatagaattattctg  
taataaaaagtgcgaaaaaattgtgtctctccccccttaataataaattctatcccaaaatctacaaatgttctgtgtacacttcttctgcttmttacttctg  
ataaaatttttgaacatcatagaaaaaacggcacacaaaatactttatcatatgtttacggttmatgaccgcaatttttcttctgacgtctgggc  
ctctcatgacgtcacaatcatgtctatcgtgaaaaagtttggagtttttggaaatttcaatcaagtgaaagttatgaatttttctgtcttcttctgt  
gggggttcccttattgttctgaaggttctgaggacggcgttctgtctaaatacacaagttatgtatgacacgtatgcaagaaagatcgggaagag  
ttgggttgagggtcagtggaaggtgagtagaagttgataattgaaagtgagtagtctatgtgggttttgcctaaatgacagaatacttcccaat  
ataccaacataactgttcttactatgtcggcgttaccggccttctgtctcgcgttctgggtgacgtgaaaacctctgacacatgacgtctcccg  
gagacggtcagcagctgtctgtatgacgggatgcccgggagcagacaagcccgtagggcgcgtcagcgggtgttggcgggtgtcggggctggcctta  
actatgcccgcacagcagatgttactgagaggtgacacatgtcgggtgtgaaataccgcacagatgctgaaggagaataaccgcacagcggc  
cttaaggggcctcgtgacacgcttmttataggttaattgtatgataataatgttcttagacgtcaggtggcacatttctggggaatgtgcggcggaacc  
cctattgtttatttttaataacattcaaatatgtatccgtctatgagacaataaccctgataaatgttcaataatgtgaaaaagggaagatgtatgtat  
caacatttccgtgtcgccttattcttcttcttctgcccatttgccttctgttcttctcaccagaacacgtgtgtgaaagttaaagatgtcgaagatcagtt  
gggtgcacgagtggtgtacatcgaactgtgattcacaacagcggttaagatccttctgagagtttctgcccgaagaacgttttccaatgtatgagcattttaa  
gttctgtatgttggcgggtatattccgttattgacgcccggcaagagcaactcggctcgcgcatacactatttctgaatgacttctgtgtgactcac  
cagtcacagaaaaagcatcttaccgagtgatgacagtaagaagaattatgcagtgctgtccataaccatgagtgataacactcggcccaacttacttctga  
caacgatcggaggaccggaaggactaaccttmttgcacaaatggggatcatgtactcgccttctgctgttgggaaccggagctgaatgaagc  
catacacaacgacgagcgtgacaccacgatgctgttagcaatggcaacacgttgcgcaaaactttaactggcgaacttacttactgtcttccggc  
aacaattaatgactgtgagggcgtgataaagttgcaggaccatttctcgcctcggccttccgctgctgtgttattgtctgataaatctggagccg  
gtgagcgttgggtctcgggtatcttgcagcacttggggccagatggttaagccctccgctatctgtatctacacgacggggagtcagggcaactat  
ggatgaacgaaatagacagatgcgtgagataggtgcttcaactgataagcattgtgtaactgtcagacaaagtttactatatactttatgattttaa  
acttcaatttttaatttaaaggatctaggtgaagatccttmttgaataatctatgaccaaatacccttaacgtgagtttcttccactgagcgtcagaccccg  
tagaaaagatcaaaaggatcttcttctgagatccttcttctgtcgcgtatctgtctgttgcacaaaacacacccgctaccagcgggtgtgtgttgcgg  
atcaagagctaccaacttcttctcgaaggttaacttggcttgcagagcgcagataccaataactgttcttctagtgtagccgtatgtagccaccactt  
caagaactctgtgacccgctacatactcgtctgtatcttctgttaccagttggctgtcctcagtggtgataagtcgttcttaccgggttggactca  
agacgatagttaccgggataaaggcgcagcgggtcgggtgacacacagccagcttggagcgaacgacctacaccgaaactga  
gatactacagcgtgagcatgagaaagcgcacgcttccgaaggagaaaggcggacaggtatccggttaagcggcagggctcggaaacaggag  
agcgcacgagggtgttccagggggaaacgctgtatctttagtctgtcgggttccgaccttctgactgagcgtctgattttgtgtatgtctgtc  
aggggggctggagcctatggaaaaacgcagcaacgcccgttmttaccggttcttggccttmttctgtgcttctcactatgttcttctgtgttacc  
ctgattctgtgataaccgtattaccgctttagtgagtgatcagctcgcgcagccgaacgaccgagcgcagcagtgatgagcagggagag  
cggaaagcgcgccaatacgcgaacggccttctcccgcggttggcggttcttactaattgagcgtggcagacaggttcccgactggaagcgggc  
agttagcgcgaacgcaattatgtgagtttagctcactcattagcccccaggttcttacttctgcttctgctgtatgtgtgtggaattgtgagcgg  
atacaatttcacacaggaacagctatgacatgattacgccaagctgtagmtaacaatgacttactaactatttctttaaattttcagagcct  
aaaaatggctgaaatcactcacaacgatggatagcctaacaacttggaaatgaaat



SEQUENCE ID NO. 6

[illegible]

pGN110:

SEQUENCE ID NO. 7

gatccctcaaaatcgtctccgctctgaaaaacgaaagtgaccccttgacatccgaaaaaatggcgaaaaaatgaaatgagccttttgggtcgaaaa  
aaatgnttttagaatctgagaaacggttaaacacgaaagatcataatttttagagaccggatgctctgaaaaatgctgacatagatttaaaaaagcata  
atatatttccattttcaacgtgaaaggtttgtgcaactttatagaatctcttattggcacatgttttttatttaactgaggcagtttgaacaccccttttgaac  
tgaatctctttgaagtatactgtcgaagagactgactgagcgttcgaaatgccagaagaaaactataattgaaatctcgcgttaattgaaatgcaacc  
cgctccacttggacaattggaaaaaaattttatcgaggcgacacccggtattttcgaatgattttctgtattttctcattttataaattcttcttgant  
atcgtctgtttgtgaaatmaantattcaaacctttttatagtagatccgggtgtaccgctagccgtacgaacccgggattggccaaaggaccca  
aaggatgtttcgaatgatacataacataagacattttcaggaggacccttgccttggagggtaccggatgactgctccaaagaaaggcgttaagc  
tcattgaacacgalttaacatcgctaagaacgacttcttgcacatcgaactggctgtatcccggttaacacatctggctgaccattacgggtgagcgtttagct  
cgcgaaacagttggcccttggagcatgagctttacgagatgggtgaagcagcgttcgcgaagatgttggagcgtcaacttaagctgggtgaggttgcgga  
taacgctgcccgaacgctctcactactacccctactccctaagatgattgcacgcatcaacgactggttggaggagtgaaagctaaagcgcgggcaag  
cgcccgacagccttccagttctgcaagaaatcaagccggaagccgtagcgtacatcaccattaaagaccacttggcttgccttaacacgctgaca  
atacaaccgttcaggctgtgcaagcgaatcggtcgggcccattgaggacgggtcgttcggtcgtatccggtgacctgaagctaaagcacttcaa  
gaaaaacgttggaggaaacacacgaacgctgtagggcacgtctacaaagaaagcattatgcaagttgctgaggctgacatgctcttaagggtcta  
ctcgggtgagcggcgttgccttctggtcgtataggaagactatctatgtaggagtagcgtcgtcagatgctcattgagtagcaaccggaatggttag  
cttacaccgccaaaaatgctggtgtagttagtcaagactctgagactatgaactgcacctgaatacgtgaggctatcgcaaccggtgaggtgcg  
ctggtctgcatctcctgatttccaaacttgcgtgatttctcttaagccgtggtgactggcattactgggtgtgtgtattgggttaacggtcgtcgtccct  
ggcgtgtgtgctgactacagatgaagaaagcactgagctgctgacgaagacgttcatgctgaggtgtacaaagcgttaacattgctgcaaaacacc  
gcatgaaaaatcaaacgaagaaagctcctagcgttgcacacgtaatacacaagtggaagcattgtccggtgagggacatccctgcgattgagcgtgaa  
gaactcccgatgaacccggaagacatcgacatgaatcttggagctctacgcgtggaacgtgctgctgctgtgttaccgcaagacaaggctc  
gcaagctcgcgtatcagccttggattcatgcttggagaagccaataagtttgaaccataaggccatctgttcccttaacattgagctggcgcg  
gttcgtgtttacgtctgtlcaatgttcaacccgcaaggttaacgatatgacaaaggagctcttaccgtggtgcaaggttaacccaattggaaggaag  
ttactactgtctgaaaaatccaggttgcacactgtgcgggtgtcgaataaggttcttctgagcgcgtatcaagttcattgaggaaacacaggaacat  
catggtcttgcgttaagctcctcactgagaaacattgtggtgctgagcaagattctcgttctgcttctgcttctgcttctgagtagcgtgggtacagc  
accacggcctgagctataactgtctcccttccgttgcgttggacgggtcttctgctgcatccagcacttctccgcgtgctcctgagatgaggtggtg  
gtcgcgcgttgaactgttctctagtgaaacccgttcaggacatctacgggattgttctgaagaaagtcacagagancgtgcaagcagacgcaatcaatg  
ggaccgataacgaagtagttaccgtgacccgtagaagacactggtgaaatctctgagaagaaagtcaggcactaaggcactgtgctgtcaatggct  
ggcttaccgtgttactcgcagtgactaaagcgttcagtcagcgttgccttacgggttccaaagagttcgttccgtcacaaggtctggaagatac  
cattcagccagctattgatttccggcaagggtctgatttccactcagccgaatcagcgtctgctgatacatgctaaagctgatttgggaatccgtgagcgt  
gacggtgtgagctgctggttgaagcaatgactggttgaagctgtctgctgaagctgctgaggtcaagataagaagactggagagattcttc  
gcaagcgttgcgtgtgcttgggttaactctgtagtcttccgtgtgtgaggaatacaagaagcccttaccagacgcgttgaacccgtgattcttctcgtg  
tcagttccgttaccagccttaccattaacacaaacaaagatagcgagattgtagcacacaacagagagctgtgtatcgtctccttaactttgtacacagcca  
agacgttgcacacgttgcgaagactgtgtagtgggtgacacgagaagcgaatcgaatcttctgacgtattcagcactcttccgttaccatttccggc  
tgacgtctcgaacctgttcaagcagtgccgcaaaactatgttgacacattagcttctgtatgtactgtgctgatttctacgaccagttcgtgaccag  
ttgcacgagttcattgacaaaatgccaagcacttccggttaaggttaacttgaacctccgtgacatcttagagtcggacttccggttgcgttaagggt  
ccctctgtcgtgagtcacaaatcactgaaactccaaaggcagccagtgaggaaagcgtgaagaaagaaagaaagagtcactgaacaggttgaattt  
cttctgtgcaaaaagatgaaatattgaattttcagccagatactccaaactagcagcgagaaagctcgaagctgttccagagtcgcccagagaatcgc  
gggaaagttagccaaagaggtatgttttcaaaaatcaataactgacataattttattgttggtaatttaagaaaataattcgaaaattccttgaattat  
caagattgcagttattttcagaaaaattgagattttcatagagctattgttaatttcttgaatttcagactgaaacttccgaaaaatcaagagaaaaatcaa  
agaaaaagtagcgggagtagcagcctggcacaccgaacagctatagaagccgggaaacttccagcctccaaaaaggttccaaaggagacag  
gtttgtcaaaagcttctcgtgattatttcaattttcagagaatcagagcttccgtgaaaaatccccgggttctcaagatctcccaagaggttca  
gcagcttcccgtcagatctcttagacggcgccgagaagagctcagaagaaagcaatccgaagagccagcaccgtaccagagaaaaaga  
agaaagagccgctgagatatttctagcaacaagaaccggaggagcattatctacccaccgccaaacttgcactatgcaacaacagatttagtgataagca  
aagtgaacagtagtagaattggaaggaatgaaagaaagatttcacggattgttgaacagagtcacacgcaagaatctgttcaattgtca  
gagaacttcaagagaatgtgattcgttcaaggtgagtagaatacgaaggaagaaatattttttcaggggacttcttgcgtg  
acattttcaagctcagccttccaccaggttcttctaacgtctatgacgttggcggcagttatcaactcgaatttccctatgtcgtgtaacttctc  
cgtctcgtgattgacagttcaaaagaggttccgtgaaatgacagagggcgtacgggtgaacgtgattcaattatcgcacatttgaattatcaacaag  
tgcctcagcaaggttcttgcgtgaaatcattctgagcttgaagaaccaactgattgattcagttgaggtcgcatttgcgttccgtgaaagaggttgcga  
gcaaaagcttctgagagattgctcagcagcttcaacagttctacgaccgttctgtgcaattctcattggaactgaaagatcggaaaatgactggatc  
gacgtattcagttatgattgagactgcaatgcagattcgaaggaacaaattgctgtaaggtagaataataaattagtttagaaaaataaattag  
aataatmaaatctctactagccaatcagggcacccttttgcgcatgttcttattgaataatttggagaatttctatctcgtcggaaatctggaatt  
cgacgagatcttctgcttctgtcagctgcacgcttctgtctcccttctcgttcttctgtgtacaccaagaaccttgttgaattcactgaattc  
gtgactgcttctgtctcagtgatgactagacgactgattctgagaacacagattgaggtgcgattgaggtgacctgaattggaggaataatcagaa  
cttttgaataatcagcaggaattaaaaaattttctcagacattctcaaaaatttactgaccaltgttgcctcaacatttttataaagttatgaaaa  
aagtgtgaaaaatcggaaaattggcaatttcagaccatttttaagcattttcaaaaaaaattgcagctgaaataaagtctattttcagataaaatcagcgaat  
ttctgttctgtgacacatgttttagtttttaaaaatgttggagaacatgtgtgcaataggttaatttcatagaatttcattgtttttttcaatcaacaaatc  
caaatcttcaaaactcacttttgcggagctgggtatcaagaatcgtcagtttttaagacgagcattctgtatcactgaaaattttttatca  
aaacttgaataatcaactaaaccacatttaacattctcgtctgtctggttgcgtatgacgggtgaagaaagccaaattgtagtagttgattgggttcaagt

[illegible]



pAS2-cyh2-HA+bothT7-final

[illegible][illegible]

gggtaacgccagggtttccagtcacgacgttgttaaacgacggccagtcgtccaagcttccgagctcggagatcccgagctmgcaaataaagccttc  
gagcgtcccaaaacctctcaagcaaggtttcagatataatgttaacatgcgtacacgcgtctgtacagaaaaaaagaaaaattggaatabaataaacgttct  
aatactaacaataactataaaaaaataaataagggaacctagacgttcagggtgtcctaactctctcttttcgggttagagcgagatgtgggggagggcgctgaigt  
cgctgacataactaattacatgatactctttgtgtttccggggtgtacaatatggaacttccctcttttcggcaaccaaaccatacatcgggattcctataactct  
ctgttggtctcctaactatgttaggtgtggggggagatatacaatagacagacatcagacaagagataatgggtctaaacagactacaacaaactacacgt  
ccctatgtatgtgtgtgtacatacgaacgaataactctgtacccctagccttagatcgaatcactatcagctcgaatgtaactcctctctctctctctctctct  
ataataggcgcatcgaact  
aaattaacgacaagaagacagaccacaagatgtcgtgttccagagcgtatgaggggatctctcgaacacacgaaacttttctctctctctctctctctct  
ctctctaatgagcaacggtatcaggcctctctccagttcgtgaattgaaataaaaaaagttgcctctctctctctctctctctctctctctctctctctct  
ttgtttct  
gaaagatgaagctact  
agaacaactgtggaggtctcgtct  
gctattctactgtattmccctgagaagaccttgacatgttgtaaaatgagatcttctacagatataaagaacgattgacataaagacataatgtgt  
aataaagatgcgct  
agtaacaaagggtcaaaagacagttagctgtatcgcgggaattcttaatacgaactcactataggcatatggccataggccatggaggcccccggg

[illegible]





pGN205:

SEQUENCE ID NO. 10 -

ccggtgttacggggccccccctcagggtcagcgtatcagataagcttcgattgaaaaagagataaatggacgatgggaagagctctcgt  
gttccaggagatcagaaaaagcaactgttccaaatcctaaggaggagagaataatcaattcagaatttctgctcgaacaggctgggaactggaga  
tcttctgtatcttctgtatgtgtgtgagcgaagcgaagaaacctgtctcagaatcattcgtgaagatcttctgatacaactgtcaaggctggagccac  
tctcaaggtcattgttcatattgtgtgtgagccagcaccagatgtacatgtgtcattcaatggaaaaggaaatcgagagagcaaggctcaaaattgaaa  
tgagccatacatctcagatgt  
actatcaatatacaggtaaaaagcaagcgaacgaacaaagggaacaaatcgaggtaactgtatgtcttgaagatcgtgtcaactctgtactggaac  
caccagaggatgacggaggagagcgaattgagttctatgaattgaaaagatgaacaccaaggacggaatctgggttcctgtgtgacgtatggag  
atacccacttcacagtcgattcactcaacaaggagatcattacaagtctcgtgtcaggctgtcgaacagcgaagacaccttctgtatcattggaactg  
aaacgataatttggctaaaaatccattgtatgt  
ccactagtctagaagcgtgtcgaagggt  
agaaaaagagtcattcgaacaggt  
gcaagtcgttcacagtcgtccagagaaatcgagggaagtgagccaagaggtatgttttcaaaaaataaactgtatcataattttatgtgtgtgtgtgt  
aagaaaaataatttgaaaaatccttgaattatcaagattgcagtaataatttcgagaaaaattgagatattcatagagctattgttaatttctgtgtgtgt  
actgaaactctggaaaaatcaagagaaaatcaagaaaaggtatgacgggt  
accagctcctcaaaaagggtcgaaggagaccaggt  
ccggtgtgtcgaagatctccagaaagggt  
aagagccagcaccgctaccagagaaaaagaaagagcgggt  
cgactatgtcaacaagattgtgataagcgaagtgacagatcagagaatgaattgggaagaaatgaagaaaaagatcagcgaattgtgtgtgtgtgtgt  
agtcacagcgaagaaatctgtcaaaattgtcagagaaatcttctcagaagaaatgt  
taatttaattttcagagggtcttctgt  
tcgaaattcctcattgt  
caaaatcctcgcacattgtatcaacaagggt  
agtcgccattgt  
gaaactgaaagatcgaaaaatgacgt  
tatataaatgt  
aaatttcataattctgt  
ccaagaacctgt  
aggggtgacctgaaatgggaataatcgaacttttgaataatttcaggaggtataaaaaattttctgacaatctcaaaaatttactttatgtgtgtgtgtgt  
tgcttccaaatmttcaataaaaggtatgaaaaaatgtgaaaaatcggaatgtgcaatttcagaccatttttaagcattmcaaaaaaattgtgagctga  
aataaatgtcatttcagataaatcgagcgtatttctgt  
aattmccatgt  
atctctgtatcactgaaaaatmtttaaatacaaaactgtataatcaactaaaccacttataactttcagatcttctgtgtgtgtgtgtgtgtgtgtgtgt  
gaagccaattgt  
aagtgtatataattgt  
accaaatcgataatccatgt  
caaaaaataggggttttaagtggt  
ttggagcagcttctgt  
catcgaagctgtcttaaaaaatgtcattaaaaaggggt  
tcacaacaataaaaatgt  
gtaaatttcagttcaattttcattttacacataatcataaagctacgt  
caaaaagaacaagaagaacaagcagctgt  
gagggggagactgaagatggcaattgaggacagtgctgt  
ctaagcccttctgt  
aaataactgaagaagaataaataattgt  
cttaatttggagagatgt  
gttcaaaactagatccagaattcgaagaacagagaggtttatgaggagatccgtgaagaaatcattggaacgcggatatttcgagatgagatgtgtgt  
gcgacagtggt  
tcagaaggtatcattttgagtttggcccgcaaatctgt  
aaaaacttttgt  
aaaaatgaagattccagacagcatgaggtcagcgt  
gttt  
cgcgatgt  
tactttgaaaagcttctcaggacacgtattccagattcaccgaattgacatcaaaaactcgggaatttggctgtgtgtgtgtgtgtgtgtgtgtgtgtgtgt  
gatgt  
gagcgatgt  
cattgt  
gtgaatgt  
gactcgt

bioRxiv preprint doi: <https://doi.org/10.1101/2020.03.10.332000>; this version posted March 10, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

[illegible]

pGN207:

SEQUENCE ID NO. 11

ccgggtggtaccgctgacccgaacccggggtctagaactagtggtatccacttgagatcaacatgatcagaatcccaatcagttggctctggacga  
cctggtctatctgacgaatcgaatggatmragccaaaatacgggttcacgttccaatgatcagaaggtcctctgctgttgacagccttgacacggaa  
ctgtgaatgatctccctgtgagtgaaatgactgtgagtggtgtatccactacgtccacatggaacccagatcctgctgtgttcatctttcaatf  
catagaactcaatngctctccctgctatcctctgtgtgttccagtaagatgtgcacgatcttccgaagacatcagttacatcgtgtgctccctgtgt  
cgttggctgtcttttaccctgataatgatagtgacactgtcagttccattaatgtgtgtgagtgatggtatattmccacttgcctacgaagtgcttggca  
aagcaaatctcagatgtatggtcctatctcaatmgagccttgcctcctcgattcttccatggaatgaccatgttaccatctgtgtgctcaccatca  
atatgaacaatgaactgagagtggtcctcgaccttgacagtggtacgaagaatcttccagatgaattcttgagcaaggttctgtgtcgaacaac  
acgatcagaaggatcagaaggatcctcagcttgcagccttgcagcagaatctgaattgatacttctccctctaaagatttgacagtggtgt  
ttctgatctctggaacaacgagagcttctccatctgtccttcttcaatgacgaagcttatcgataccgtcgaactcgtgagggggggc  
cctcgtcgtgagtggtcacaatcactgaaactcgaagggcagcagtgaggaacgtgaagaaagaaagaaagagatctgaacaggttgatmtct  
ttctgtcgaagaaatgaatnatgatttccagccagatactccaaaactagcagcgaagagctgtcgaagctgttcacagctgcccagagaatcgcg  
ggagtgagccaaagaggtgtgttttcaaaaatcaataactgataatattttatgtgtgtgaatttaagaaataatctgaaaatctctgaaatc  
aagattgcagatattatcagaaaaattgagataattcatagagctattgttaatttctgtattcagactgaaactcggaaaaatcaagagaaatcaaa  
gaaaaggatgacggggatgacgctgtgcacccgaacagctatagaagccgggaaacttaccagctccaaaagggtccaaggagaccaggt  
ttctcaaaagcttctcgtgattatctcattmcaattttcagagaatcagagcttctgaaaatcccccgttctgacgaatctccagaaggttctcag  
cagcttccccgtcagcagatctcctagacggcgccgagaagaaagctcagaagaaagcaatccgaagagccagcaccgtaccagagaagaaagaa  
gaagagcgcgtgtgataattctcgaacaagaagctggagagcatatattccaccgcccaactcgaactatgcaacaacagattatgataagcaa  
agtgaacagatcagagaatgaatgggaaagaaatgaagaaaaagattcagcagtggttaacagagtcacgcgaagaaatctgttcaaatgtcag  
agaactttcagaagaatgtgattcgttcaagtgagtgagaaaaatgaagaaaaagaaagaataatattttcaggggacttctcgtcgtgac  
atattcaagctcaggttctcaccaggttcttcaactgtatgtcagcttggcggcagttatcaactcgaatctcctatgtcgtgagacttcttcc  
gtcgtctgtgtgacagttcaaaagaaatctcgtgagaatgacagagggcgtcaggtgaacgtgatcaaatctatcgcacattgtatattcaacaggt  
gtcaggaaggttctgctgtggaatcagatctgtgaggaacaaactgtatgattcagtgaggtcgtccattgtcgtgaaagagtggtgag  
caaaagcttctgagattgtctccagcagcttcaacagtgctacgaccttctgtgcaatctcatggaactgaagatcgggaaatgactggtatc  
acgtattcagatattgattgagactcgaatgcagatcgaagggacaaatttgcgttaagtgataataataatgtttatagaaaaaataataga  
ataattaaatctctacagcccaactggtcgtcagctatgttatttgaagaaatgtgagaatttctcatattctcgtcgggaaatctggaattc  
gacgagatcttctgcttctgtcagctgtcagcttctccttctcgttctcgtgtgtacaccaagaacctgtgtgagttcattcactgaactgt  
gactggtctgtgtcactggtgactacagcagctgattctcgaatcagatgagtgctgattaggggtgacctagaatgggaaataatcgaact  
ttgaaaaatctcagggaggaataaaaaatattctcgaactctcaaatatttatttgcacattgtgtcctcaacattttcattaaagttaataa  
atgtagaataatcggaaatgtcaattttcagaccatttttaagcattmcaaaaaaaatgtcagctgaataatgtcattttcagataaatcagc  
ctgtgtgtgacactagttttatgatttttaaaatgttgaagaacatggtgcaataggttaattcagaaatcctgtgttttttcaatcaacat  
aaatcttcaaaactcacttttgcggagctgggctatcaagaatctgtcagttttatagacgagcatctctgtatcactgaaaatattttatca  
aactgaaatatacaataaaccacttatttaacmctcgaatcttctgtcgttctgtacgaggtggaagccaaatgtgtgtgtgtgtgttcaagtc  
cttctggt  
gatctttgt  
ctaggaaactgt  
tagctaaacggzaaattttatgttttttttcaagaaccactgaaacccctaatgtatataattttggagcagcttctgtgtgtgtgtgtgtgtgt  
tcgataaaacagaatttaagtgttaattgttcaattttatcaaatgt  
aggggtttcagtggtttttcattaaaaagcttaatttaactaaaaatcattatmtccaaactgtgtcacaacataaaatgtgtgtgtgtgtgtgt  
aaaaaatgttt  
aatcataaagctacgtctgtatctctcgcacttacctgcgcctgattcgaagaacaacccgtatgcccagaagaagaagaacacacacacacac  
ggtagtgtgacgttctacgcgaatcgtacgaatgt  
gt  
aatccgaagaatgt  
aggcgtatccggcggtcgt  
ggcttagtgaagt  
ggaggtttatgt  
gatgtggaagaggtctcgaagaagactacagagattttatatactgtatcagaatgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgt  
atcgtgaattgt  
ttcagagaagcttaccgacaaatgcaatcattmgtgactcaagaagctgtcacaatgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgt  
gagtgtgcaagaagaaatatttgaccaaaaaaccaaccaataaaatt  
atgtatt  
cctacgagcgtatctacggaatgtctatcgaacgttctgtccgacttgcctgaataccagcaatactmgaagaagctgtgtgtgtgtgtgtgtgt  
gattcaccgaatgacatcaaaaactcggaaattgt  
aatgaccgaagaggaacacacttcttggcagaatctataataataatatttgaactgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgt  
ttactgtcgt  
aacagcgcagcatttgcacacncttcaaatgattgt  
agggaatgt  
gactgtcttgcagttctcattcttctcagaatcagagccaccaggaaagaaagaaagaaagaaagaaagaaagaaagaaagaaagaaagaaagaa

bioRxiv preprint doi: <https://doi.org/10.1101/111111>; this version posted January 1, 2017. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

gaaaaagagaatattggtgcgacgggagcgtgggagacagagagctgaacgicactgtgatcaaaagtgtggagaacaaggacaaaggatcgtcgacg  
tcgccaggattctgacgaaaaatcgtcgccgaacgagagatgaccgcaaggatcggagtaaaagatcgtcgtcgtcaagactcggatgatgagg  
atcggaagggtcgtgaacgtcgcggaagatfcagggggaaagacgtcgcggagatcgggagtcgacgtgcatcgaacaaggatcaggaggatcccg  
tgaagatccgctgacccgaagcagagatcgtgaggcaicgacgtgacccgtcgtcatgactctgatgatgatcgtaaaaactgtcgggagataaagt  
gaaagcgcgagggagacgtcgtcgtgaagtgaatcgtgatcgaacccgacgtcgttgaatttcaaatmtaataactgataattgttttttctatt  
aattattattctcttgggtgtttttttctgtttcaaaaaaattcaaatcaatcaaatcaaatgagcgtgtttttttcttccgtctccaaatcgtattccgt  
ccctcactcgaacacaatgtcgaagtgtttattattcttctgtctttactattaggagagtggtgggaagtgtgggaagggtgggaacacacaaaagcgtg  
atggaaatgaaataaggacacacaataatgcaacaacattcaatcagaataatgaggaaggttttaaaagaaaaacataaaatataataggaggaa  
ggaaaactagtaaaaaataaggcaaaagaaattaggcgaacgatgagaattgtctcgtcgttggcaaatcggaatccgtatggagaggcacgttttggcg  
aggcaaatgttctggfatggagatcgttaaaaaatttaagtgaaaattgtgttgccttttcaaaaatttccgattttcgttgaataacggtgccaggctc  
cgacacgtcttccaaattttcaaatcaaaagagccmttaattggcgtgtagtgtcctaatttctcgttttgaaaaatttttccgttttaacgaaattgtatgtatt  
tattatgattmcaataaaattcaaaagaaactgggtgaaaactcgggaaaattgtgaactacagtaatccaatcttaaaggcgcacacctmtaaatgtccgc  
cccaactcgaatattmttaagattccttgatgagcggcgcccgccaggcgtggagctccaaatcgccttatgtgagtcgtattacaattcactgtgccgtcgt  
ttacaacgtctgtgactggggaacacccgtcgccgttgcacccaacttaactcgccttgcacatcccccctgccagctggcgtaatatgcgaaggcgcc  
gcaccgctcgccttcccaacagttgtcgtgactgagtgatgcgaatggcgcgcgcctgtgtcggcgcgacttaagcgcgcggcgccgtgttgaccg  
gcagcgtgacccgtacactgtccagcgccttagcgcgcgccttctcgttctcccttcccttccgtccacgttgcggcgcttcccccgtcaagctcta  
aatcggggggctccctttaggggttccgatttagtgccttaccggcacctcgaccccaaaaaacttgattagggtgatgtgtacgtatgtggccatcgccct  
gatagacgggttttccgcttggacgttggagtcacgttctttaaagtgtgactctgttccaaaactgggaacaacactcaaccctatctcgtgttatcttmt  
gattataagggaatttgcgaattcggccctattgtgttaaaaaatgagctgatttaacaaaaaattaaacgcgaatttaacaaaaatattaacgtttacaatttca  
gggtggcacttttcggggaaaagtgcgcgggaacccctattgtttatttttcaataacatcaaatatgtatccgtcctatgagacaataacccgtgataaagc  
ttcaataatattgaaaaagcgaagagatgatgattcaacatmccgtgtcgccttattctccttttgcgcctatttgcgcatttgccttctgttttgcctcaacgaac  
cgtgtgtgaaagtgaagaagctcgaagatgttgggtgtgcacgtgtgggttatcgcgaactgactcaaacacgcggtaagactccttgagagcttgcctgc  
ccgaagaacggtttccaatgatgagcactmttaaaagtctgtatgtgcgcgggtattatccgttatgacgcggcggaagcaactcgtgtcgcgcgat  
acactattctcagaatgactgtgtgagtactaccagtcacagaaaagcatcttaccggatggcagtagacagtaagagaattatgcagtctgtccataag  
catgagtataacactgcgcgccaacttactctgacacagatcggaggaccgaaggagctaacgccttttttcaacaatgtggggatcatgttaactcg  
ccttgatcgttgggaacccggagctgaatgaagccataaccaaacgacgagcgtgacaccacgatgcctgtagcaatggcaacaacgctgtgcgcaact  
attaactgtggcaactactactctgacgttccggccaacaattaatagactgtgagaggcggaataaagtgtgacgaccacttctgcgtcgcgccttcc  
ggctgtgcgttttattgtctgataaatctggaagccgggtgagcgttgggtctcgcgggtattctattgacgacgttggggccagatgtgaagccctccgtaicgt  
agttacttaccagcgggacgtcagcaactatgtgatgagcaagaatagacagatcgtgagataggttccctcactgattgaagcattggatgaactctgag  
accaagtattactatatactttagatttttaaaactatttttaaaaggaactatgagtagagacttcttgataaactctatgacccaaaactccctaa  
cgtgagtttctgtccactgagcgtcagacccgtagaaaagatcaaaagactcttcttgagactctttttctgcgcgtgaatctgtcgtcgtcgaacaaaaa  
aaaccaccgctaccagcgggtggmttttgcgggaicaagagctaccaactcttttccgaaggtaactgtgcttcagcagagcgcagataccaaactact  
gtccttctagtgtagcgtgtatggccaccacttcaagaactctgttagcacccgectacatcctcgtctgtctaactcgtgtaccagtgtgtcgtccag  
tggcgataagctgtgtttacccgggtgtgactcaagacgatagtaccggataaggcgcagccgtcgtgtgaacgggggggttcgtgcacacagcc  
cagcttggagcgaacgacctacccgaactgagatacctacagcgtgagcattgagaagaagcggccacgcttccgaaaggagaaaagcggagacag  
tatcctgaagcgcgaggctcgaagcagagcgcagcagggagcttccagggggggaacgccttggtatctmtatgtcctgtcgggtttccgccac  
ctctagttagcgtcgtgatttctgtcgtcgtcagcggggcgcgagcctattggaaaaacgcgcagcaacgcggcctttttactcgttccgtcgtcgtcgt  
ggcctttgtctacatgttcttctcgtctatccctgattctgtgataacagctattaccgctttagtctgataactcgtcgcgcagccgaaga  
ccgagcgcagcgtgagtcgtgagcggaggaagcgggaagagcgcgaacacgcgaacccgcttccgcgcgcgttggcgcgattcaatgacgtg  
gcacgacaggtttccgactgtgaagcgggcagtgagcgcgaacgaatnaatgtgatttacttactactataggcaccgccaggcctttacactmtatgct  
tccggtcctatgtgtgtggaattgtgagcggataacaattcacacaggaacacagctatgacctgattacgccaaagctcggaaftaacctcactaa  
agggaacaaaaagctggggggggtcttccaaaaactgtcttccgctcgtgaaaaacgaaagtgcacttttgacatccgaaaaaagtggcgaaaaaatga  
aattgagccttttgggtcgaaaaaaatttttgaatgctgagaacacgftaaacacgaagaatcatattttttgagaccgggatgctcgtgaaaaatgtctg  
acatagatttttaaaaaagcatatataatttttcaatttcaacgtgaaagtgttggcaactttatgaatctcctatnggcacatgttttttaaacigtaggcag  
ttttgaacaccttttgaacattgaatcttggatatactgtcgaaaagactgtgacgttgcgcgttgcgaatgtccgaacgaaactatatttgaatctgcg  
gctaaatgagaagaatgaaacccgctcactgacgaatnggaaaaaatttttccggagcgacaaagctattttcgaaaatgtatttctgttatttct  
cattttataaattcttcttatttatcgttctgttggagaaatttaattgtattcaaacattttatagtaagata

Claims

1. A method of identifying DNA responsible for  
conferring a particular phenotype in a cell which  
5 method comprises

a) constructing a cDNA or genomic library of  
the DNA of said cell in a suitable vector in an  
orientation relative to a promoter(s) capable of  
initiating transcription of said cDNA or DNA to double  
10 stranded (ds) RNA upon binding of an appropriate  
transcription factor to said promoter(s),

b) introducing said library into one or more of  
said cells comprising said transcription factor, and

c) identifying and isolating a particular  
15 phenotype of said cell comprising said library and  
identifying the DNA or cDNA fragment from said library  
responsible for conferring said phenotype.

2. A method according to claim 1 wherein said  
20 library is organised into hierarchical pools prior to  
step b).

3. A method of assigning function to a known  
DNA sequence which method comprises

25 a) identifying a homologue(s) of said DNA  
sequence in a cell,

b) isolating the relevant DNA homologue(s) or a  
fragment thereof from said cell,

c) cloning said homologue or fragment thereof  
30 into an appropriate vector in an orientation relative  
to a suitable promoter(s) capable of initiating  
transcription of dsRNA from said DNA homologue or  
fragment upon binding of an appropriate transcription  
factor to said promoter(s),

35 d) introducing said vector into said cell from

step a) comprising said transcription factor, and  
e) identifying the phenotype of said cell  
compared to wild type.

5           4. A method according to any of claims 1 to 3  
wherein said DNA library, homologue or fragment is  
cloned in a sense and an antisense direction relative  
to said promoter.

10           5. A method according to any of claims 1 to 3  
wherein said DNA library, homologue or fragment is  
cloned between two promoters capable of producing  
dsRNA from said DNA library, homologue or fragment  
upon binding of said transcription factor to said  
15 promoters.

            6. A method according to any of claims 1 to 5  
wherein said cell is adapted to express said  
transcription factor.

20           7. A method according to any of claims 1 to 6  
wherein said DNA library, homologue or fragment is  
constructed in a suitable vector which comprises a  
sequence of nucleotides encoding said transcription  
25 factor operably linked to a suitable promoter.

            8. A method according to any of claims 1 to 6  
wherein said transcription factor is encoded by a  
further vector independent of the vector including  
30 said DNA library, DNA homologue or fragment and which  
sequence encoding said transcription factor is  
operably linked to a suitable promoter.

            9. A method according to claim 7 or 8  
35 wherein said transcription factor comprises any of T7,

T3 or SP6 polymerase.

10. A method according to claim 7 or 8 wherein  
said suitable promoter comprises any of let 858,  
5 SERCA, UL6, myo-2 or myo-3.

11. A method according to any of claims 7 to 10,  
wherein said suitable vector or said further vector  
comprises a selectable marker.

10

12. A method according to claim 11 wherein said  
selectable marker comprises a nucleotide sequence  
capable of inhibiting or preventing expression of a  
gene in said cell and which gene is responsible for  
15 conferring a known phenotype.

15

13. A method according to claim 12 wherein said  
nucleotide sequence comprises a sequence which is a  
part of or identical to said gene conferring said  
20 phenotype, and which nucleotide sequence is itself  
oriented relative to a suitable promoter(s) capable of  
initiating transcription of double stranded RNA upon  
binding of an appropriate transcription factor to said  
promoters.

20

25

14. A method according to claim 12 wherein said  
nucleotide sequence is a part of or identical to said  
gene sequence conferring said phenotype, and which  
nucleotide sequence is such as to permit integration  
30 of said suitable or further vector by homologous  
recombination in the genome of said cell and following  
said integration said nucleotide sequence is capable  
of inhibiting expression of said gene sequence  
conferring said phenotype.

30

35

15. A method according to claim 14 wherein said nucleotide sequence comprises stop codons sufficient to prevent translation of said nucleotide sequence following its integration into said genome.

5

16. A method according to any preceding claim wherein said cell is a microorganism suitable for feeding to, transforming or infecting an organism.

10

17. A method according to any of claims 1 to 14 wherein said cell is contained in an organism or an embryo thereof.

15

18. A method according to any of claims 1 to 17 wherein said promoters are T7 promoters.

20

19. A method according to any of claims 12 to 18 wherein said known gene sequence comprises a sup-35 gene or a fragment thereof which is selectable by identifying offspring growing at a temperature above 25°C following introduction of said vector in the genome of a pha-1 et123ts mutant *C. elegans* worm.

25

20. A method according to any of claims 1 to 19 wherein said cell or organism is contacted with a specified compound for screening for a desired phenotype, such as resistance or sensitivity to said compound when compared to the wild type cell or organism.

30

21. A method according to any preceding claim wherein said transcription factor is inducible.

35

22. A method according to claim 16 wherein said microorganism is an *E. coli* strain which is an RNAase



III and preferably an RNAase negative strain.

23. A method according to any of claims 17  
wherein said organism is selected from the class  
5 nematoda.

24. A method according to claim 23 wherein said  
organism is *Caenorhabditis elegans*.

10 25. A method of generating a transgenic non-  
human organism comprising an exogenous transcription  
factor and a transgene comprising a promoter operably  
linked to DNA fragment which is expressed upon binding  
of said transcription factor thereto, the method  
15 comprising

a) providing a first transgenic organism  
comprising a first construct incorporating DNA  
encoding an exogenous transcription factor and a  
second transgenic organism comprising a second  
20 construct including at least one promoter operably  
linked to a desired DNA sequence which is expressed  
upon binding of the transcription factor of said first  
transgenic organism thereto,

b) crossing said first and second transgenic  
25 organisms and selecting offspring expressing said  
desired DNA sequence.

26. A method according to claim 25 wherein said  
first and second transgenic organisms are generated by  
30 transforming said first and second constructs into  
respective microorganisms for subsequent feeding to  
the respective organism.

27. A method according to claim 25 or 26 wherein  
35 said second construct comprises said desired DNA

sequence in an orientation relative to said promoter so as to be capable of initiating transcription of said DNA to dsRNA upon binding of said transcription factor thereto.

5

28. A method according to claim 27 wherein said second construct comprises two promoters flanking said desired DNA sequence which promoters can initiate transcription of said DNA sequence to dsRNA upon binding of said transcription factor to said promoters.

10

29. A method according to claim 27 wherein said DNA sequence is provided in a sense and an antisense orientation relative to said promoter so as to produce dsRNA upon binding of the transcription factor to the promoter.

15

30. A method according to any of claims 25 to 29 wherein said second transgenic organism further comprises a reporter gene operably linked to a promoter which is capable of initiating transcription of said reporter upon binding of said transcription factor thereto.

20

25

31. A method according to any of claims 25 to 30 wherein said transcription factor comprises a polymerase.

30

32. A method according to claim 31 wherein said polymerase comprises any of T7, T3 or SP6 polymerase.

35

33. A method according to any of claims 25 to 31 wherein said promoters comprises any of T7, T3 or SP6 promoters.

34. A method according to claim 30 wherein said reporter gene comprises any of those sequence encoding Luciferase, Green Fluorescent protein,  $\beta$  galactosidase or  $\beta$ -lactamase.

5

35. A method according to any of claims 25 to 30 wherein said organism is of the species nematoda.

36. A method according to claim 35 wherein said nematoda species is *C. elegans*.

10

37. A transgenic non-human multicellular organism obtainable according to the methods of any one of claims 25 to 34.

15

38. A method of validating clones identified in yeast two hybrid vector experiments which method comprises

20

a) providing a construct including the DNA encoding the protein identified in the two hybrid vector experiment, which construct is such that said DNA is orientated relative to a promoter(s) that is capable of initiating transcription of said DNA to double stranded RNA upon binding of an appropriate transcription factor to said promoter(s),

25

b) transforming a cell comprising said transcription factor with said construct, and

c) identifying a phenotypic change in said cell or organism when compared to a wild type.

30

39. A method according to claim 38 wherein said DNA sequence is provided between two promoters capable of initiating transcription of the DNA sequence to dsRNA upon binding of the transcription factor to

35

said promoters.

40. A method according to claim 38 wherein said DNA is provided in a sense and an antisense orientation relative to said promoter such that binding of the transcription factor to said promoter initiates transcription of dsRNA from said DNA.

41. A method according to any of claims 38 to 40 wherein said transcription factor is inducible in said cell.

42. A method according to any of claims 38 to 41 wherein said promoter is a phage polymerase promoter and said transcription factor is a RNA polymerase.

43. A method according to claim 42 wherein said polymerase is any of T7 RNA polymerase, T3 RNA polymerase or SP6 RNA polymerase.

44. A method according to claim 43 wherein said promoters comprise any of T7, T3 or SP6 promoter

45. A method according to any of claims 38 to 44 wherein said construct is such that it may be used in yeast two hybrid experiments.

46. A method according to any of claims 38 to 45 wherein said cell is an *E. coli* cell.

47. A method according to any of claims 38 to 45 wherein said cell is part of an organism or an embryo thereof.

48. A method according to claim 47 wherein said

organism is of the species nematoda and preferably *C. elegans*.

49. Plasmid pGN1 as illustrated in Figure 1.

50. Plasmid pGN100 as illustrated in Figure 2.

51. The yeast two hybrid vector plasmid illustrated in any of Figures 4, 15 or 16 (Seq ID Nos 8 and 9).

52. A plasmid as illustrated in Figure 7.

53. A plasmid as illustrated in Figure 8.

54. A method of alleviating infestation of plant pests, which method comprises

a) identifying a DNA sequence from said pest which is critical for its survival, growth, proliferation,

b) cloning said sequence from step a) or a fragment thereof in a suitable vector in an orientation relative to promoter(s) such that said promoter(s) is capable of initiating transcription of said DNA sequence to RNA or dsRNA upon binding of an appropriate transcription factor to said promoter(s), and

c) introducing said vector into the plant.

55. A method according to claim 54 wherein said DNA sequence is provided between two promoters such that binding of the transcription factor to the promoters results in transcription of the DNA to dsRNA.

56. A method according to claim 54 wherein said DNA sequence is provided in a sense and an antisense orientation relative to said promoter such that binding of the transcription factor to the promoter results in transcription of the DNA to dsRNA.

57. A method according to claim 54 wherein said pest is a nematode worm.

58. A method according to claim 57 wherein said nematode comprises any of *Tylenchulus* ssp., *Radopholus* ssp., *Rhadinaphelenchus* ssp., *Heterodera* ssp., *Rotylenchulus* ssp., *Pratylenchus* ssp., *Belonolaimus* ssp., *Canjanus* ssp., *Meloidogyne* ssp., *Globodera* ssp., *Nacobbus* ssp., *Ditylenchus* ssp., *Aphelenchoides* ssp., *Hirschmenniella* ssp., *Anguina* ssp., *Hoplolaimus* ssp., *Heliotylenchus* ssp., *Cricone mellasp.*, *Xiphinemasp.*, *Longidorus* ssp., *Trichodorus* ssp., *Paratrichodorus* ssp., *Aphelenchs* ssp.

59. A method according to claim 55 wherein said DNA sequence or fragment thereof is cloned between two tissue, preferably root specific promoters.

60. An expression vector for use in a method according to any proceeding claim comprising a promoter or promoters oriented relative to a DNA sequence such that they are capable of initiating transcription of said DNA sequence to double stranded RNA upon binding of an appropriate transcription factor to said promoter or promoters.

61. An expression vector according to claim 60 comprising two identical promoters flanking the DNA sequence.

62. An expression vector according to claim 60 comprising said DNA sequence in a sense and an antisense orientation relative to said promoter.

5 63. An expression vector according to any of claims 60 to 62 which further comprises a nucleotide sequence encoding a selectable marker.

10 64. An expression vector according to claim 63 wherein said nucleotide sequence encoding said selectable marker is orientated relative to the promoter(s) such that transcription of the nucleotide sequence to double stranded RNA occurs upon binding of an appropriate transcription factor to said  
15 promoter(s).

20 65. An expression vector according to claim 64 wherein said nucleotide sequence encoding the selectable marker is provided between the identical promoters capable of initiating transcription of the nucleotide sequence to dsRNA upon binding of the transcription factor to the promoters.

25 66. An expression vector according to claim 64 wherein said nucleotide sequence encoding the selectable marker is provided in a sense and an antisense orientation relative to the promoter such that transcription of the nucleotide sequence to dsRNA upon binding of the transcription factor to said  
30 promoter occurs.

35 67. An expression vector according to claim 63 or 64 wherein said selectable marker comprises a nucleotide sequence encoding sup-35, for introduction into *C. elegans* having a pha-1 mutation.

68. An expression vector for expressing an appropriate transcription factor for use in a method according to any of claims 1 to 48 and 54 to 59 which vector comprises a sequence of nucleotides encoding  
5 said transcription factor operably linked to suitable expression control sequences.

69. An expression vector according to claim 68 wherein said expression control sequences include  
10 promoters which are inducible, constitutive, general or tissue specific promoters, or combinations thereof.

70. An expression vector according to any of claims 68 to 69 wherein said transcription factor  
15 comprises a phage polymerase, and preferably T7 RNA polymerase.

71. An organism or cell transformed or transfected with a plasmid according to any of claims  
20 49 to 53 or an expression vector according to any of claims 60 to 70.

72. An organism according to claim 71, which is of the species nematoda and preferentially *C. elegans*.

25  
73. A method of introducing dsRNA or DNA capable of producing dsRNA into an organism which method comprises feeding said organism with a suitable microorganism comprising an expression vector  
30 according to any of claims 60 to 67 or feeding said organism directly with an expression vector according to any of claims 60 to 67.

74. A method according to claim 73 wherein said  
35 microorganism or said organism is adapted to express



said transcription factor.

75. A method according to claim 74 wherein  
either said microorganism or said organism comprises  
5 an expressing vector according to any of claims 66 to  
69.

76. A method according to any of claims 73 to 75  
wherein said organism is *C. elegans* and said  
10 microorganism is *E. coli*.

77. A method according to claim 76 wherein said  
*E. coli* strain is an RNAaseIII negative strain.

78. A method according to any of claims 73 to 75  
wherein said organism is a *C. elegans* nuc-1 mutant.

79. A selection system for identifying  
transformation of a cell or organism with a vector  
20 according to claims 60 to 63 which system comprises a  
vector according to claims 60 to 63 and said  
selectable marker comprises a nucleotide sequence  
capable of inhibiting or preventing expression of a  
gene in said cell or organism which gene is  
25 responsible for conferring a known phenotype.

80. A selection system according to claim 79  
wherein said nucleotide sequence comprises a sequence  
which is a part of or identical to said gene  
30 conferring said known phenotype, and which nucleotide  
sequence is itself located between two identical  
promoters capable of initiating transcription of  
double stranded RNA upon binding of an appropriate  
transcription factor to said promoters.

81. A selection system according to claim 79 wherein said nucleotide sequence is a part of or identical to said gene sequence which confers a known phenotype on said cell or organism, and which nucleotide sequence permits integration of said vector by homologous recombination in the chromosome of said cell or organism and following said integration said sequence inhibits expression of said gene sequence conferring said known phenotype.

82. A selection system according to claim 81 wherein said nucleotide sequence comprises stop codons sufficient to prevent translation of said nucleotide sequence following integration into said genome.

83. A selection system according to claim 79 wherein said known gene sequence comprises a sup-35 gene or a fragment thereof which is selectable by identifying offspring growing at a temperature above 25°C following introduction of said vector in a pha-1 etl23ts mutant *C. elegans* worm.

84. A method according to claim 74 wherein said transcription factor is T7 RNA polymerase.

85. A method of assigning function to a DNA sequence of a multicellular organism which method comprises:

a) providing:

(i) a first construct comprising said DNA sequence cloned in a sense direction under the regulation of a suitable promoter

(ii) a second construct comprising said DNA sequence cloned in an anti-sense direction under the regulation of the same promoter as in step (i),

in a multicellular organism which is adapted to initiate transcription of said DNA fragment from said promoter,

- 5           b) identifying the phenotype of said multicellular organism compared to wild type.

86. A method of assigning function to a DNA sequence of a multicellular organism which method comprises:

- 10           a) providing  
            i) a construct comprising said DNA fragment cloned between two promoters capable of promoting transcription in said multicellular organism,  
               in a multicellular organism capable of  
15           initiating transcription from said promoter;  
            b) identifying the phenotype of said multicellular organism compared to wild type.

- 20           87. A method for expressing a gene, cDNA or DNA fragment in *C. elegans* which method comprises :  
            i) providing a transgenic *C. elegans* expressing an exogenous transcription factor,  
            ii) cloning said gene or cDNA or DNA fragment into an appropriate vector operably linked to a  
25           promoter capable of initiating transcription of said gene, cDNA or DNA fragment upon binding of said exogenous transcription factor thereto,  
            iii) introducing said vector into said transgenic *C. elegans* from step i).

30

88. A method according to claim 87 wherein said exogenous transcription factor is a phage polymerase.

- 35           89. A method according to claim 88 wherein

5

ABSTRACT

CHARACTERISATION OF GENE FUNCTION  
USING DOUBLE STRANDED RNA INHIBITION

5

There is provided a method of identifying DNA responsible for conferring a particular phenotype in a cell which method comprises a) constructing a cDNA or genomic library of the DNA of said cell in a suitable vector in an orientation relative to a promoter(s) capable of initiating transcription of said cDNA or DNA to double stranded (ds) RNA upon binding of an appropriate transcription factor to said promoter(s), b) introducing said library into one or more of said cells comprising said transcription factor, and c) identifying and isolating a particular phenotype of said cell comprising said library and identifying the DNA or cDNA fragment from said library responsible for conferring said phenotype. Using this technique it is also possible to assign function to a known DNA sequence by a) identifying a homologue(s) of said DNA sequence in a cell, b) isolating the relevant DNA homologue(s) or a fragment thereof from said cell, c) cloning said homologue or fragment thereof into an appropriate vector in an orientation relative to a suitable promoter(s) capable of initiating transcription of dsRNA from said DNA homologue or fragment upon binding of an appropriate transcription factor to said promoter(s) and d) introducing said vector into said cell from step a) comprising said transcription factor.

10

15

20

25

30

Figure 1

caccggctccagattatcagcaataaaccagccagccggaaggccgagcgagcagaagtggctcctgcaactttatccgctcc  
 atccaggtctataattgttccgggaagctagagtaagtagttccgagttatagtttgcgcaacgttggcattgctacaggca  
 tcgtgggtgtcacgctcgtcgttgggtatggcttcattcagctccggttcccaacgatcaaggcgagttacatgatcccccattgtgtg  
 caaaaaagcgggttagctccttcggtcctccgatcgttgtcagaagtaagtggccgagtggtatcactcatggttatggcagcact  
 gcataattcttactgtcatgccatccgtaagatgcttttctgtgactgggtgagtactcaaccaagtcaattctgagaataccgcccc  
 ggcgacccgagttgctcttgcggcggtcaatacgggataatagtgtatgacatagcagaactttaaaagtgtcatcattggaaaa  
 cggttcggggcgaaaaacttcaaggatcttaccgctgttgagatccagttcgatgtaacccactcgtgcacccaactgatcttca  
 gcatcttttacttaccagcgttctgggtgagcaaaaacaggaaggcaaatgccgcaaaaaagggaataaggcgacacg  
 gaaatgttgataactcatacttcccttttcaatattatgaagcattatcagggttattgtctcatgagcggatataatgtat  
 ttgaaaaataaacaataagggggtccgcgcacattccccgaaaagtgccacctgacgtctaaagaaaccattattatcatgacatt  
 aacctataaaaataggcgatcacgagggcccttctgtctcgcggttccggtgatgacgggtgaaaacctctgacacatgcagctcc  
 cggagacggtcacagctgtgtgaagcggatgccgggagcagacaagcccgtagggcgctcagcgggtgtggcggt  
 gtccgggctgggttaactatggcgatcagagcagattgtactga

Figure 1

# PGN100

ctagcatgaacacgattaacatcgtctaaagacgacttctctgacatcgaactggctgtatcccgttcaacactctggctgaccatt  
 acgggtgagcgttttagctcgcaacagttggcccttgagcatgagcttaccgagatgggtgaagcacgcttccgcaagatggttga  
 gcgtcaacttaaaagctggtgaggttgcggataacgctgccgcaagcctctatcactaccctactccctaaagatgattgacgc  
 atcaacgactggtttgaggaagtgaagctaaagcgcgcaagcgcccgacagccttccagttcctgcaagaaatcaagccgga  
 agccgtagcgtacatcacctaaagaccactctggcttgcctaaccagtgtgacaatacaaccgttcaggctgtagcaagcgca  
 atcgggtcggggccartgaggacgaggtcgttccggtcgtatccgtgaccttgaagctaaagcacttcaagaaaaacgttgaggaa  
 caactcaaaagcgcgtaggcacgttctacaagaaagcattatgcaagttgtcaggcgtgacatgcttctaaaggttactcg  
 gtggcgaggcggtgtcttctgtgcataaggaaagactctattcatgtaggagtagcgtgcatcgagatgctcattgagtcaaccgg  
 aatggttagcttacaccgcaaaaatgctggcgtagtagtcaagactctgagactatcgaactcgcaactgaatacgtgaggtct  
 atcgcaaccgtgacgttgcgtggctggcatctcctgatttccaacttgcgtagtctcctaagccgtggactggcattac  
 tgggtggtggtattgggctaacggctcgtcgtccttggcgtggtgctgactcacagtaagaaagcactgatgcgtacgaaagac  
 gttacatgcctgaggtgtacaaagcgattaacattgcgcaaaacaccgcatggaaaatcaacaagaaagtcctagcgggtcgc  
 aacgtaatcaccaagtggaaagcattgtccgggtcaggacatccctgcgattgagcgtgaagaactccgatgaaaccggaaga  
 catcgacatgaatcctgaggtctcaccgctggaacgtgctgcccgtgctgtgtaccgcaaggacagggctcgcaagcttc  
 gccgtatcagccttgagttcagcttgagcaagccaataagttgctaaccataaggccatctgggtcccttacaacatggactggc  
 gcggtcgtgttacccgtgtcaatgttcaaccgcaaggtaacgatatgaccaagagactgctacgctggcgaaaggttaac  
 caatcggtgaaggaggttactactggttgaataccaggtgcaaacgttgcgggtgtcgataaggttccgttccctgagcgc  
 caagttcattgaggaaaaccacgagaacatcatggcttgcgttaagttccactggagaacacttgggtgggtgagcaagattct  
 ccgttctgcttcccttgcgttctgctttagtacgtgggttacagcaccagcctgagctataactgtccttccgtggcgtt  
 acgggtcttctcgtgcatccagcacttctccgcgatgctccgagatgaggtgaggtgctgcgcgggttaactgttcttactgag  
 accgttcaggacatctacgggattgttctaaagaaagtcaacgagattctacaagcagacgcaatcaatgggaccgataaacgaa  
 gtattaccgtgaccgatgagaacactggtgaaatctctgagaaggtcaagctgggcaactaaaggcactgggtggtcaatggctg  
 gctcacgggtgttactcgaggtgtactaagcgttcagtcagcgtggttacgggtccaaagagttcgggtccgtcaacaagt  
 gctggaagataccattcagccagctattgattccggcaagggtccgatgctcactcagccgaatcaggtgtggtgatacatggt  
 aagctgatttgggaatctgtgagcgtgacgggtgtagctgcgggtgaagcaatgaactgggttaagtctgctgctaagctgctgg

Figure 2





ctagaaggctctagatgcattcggttgaaaatactccgggtgggtgcaaagagacgcagacggaaaatgtatctgggtctctttatt  
gtgtacactacttttccatgtaccgaatgtgagtcgccctcctttgcaacaagcagctcgaatgttctagaaaaagggtgaaaata  
gtataaafaccgttgaaaataaataccgaacaacatttgctctaattgtgaaatagaatcttcaactataatcatctcactggatc  
cccggttggtgccaaggacccaaagggtatgttcgaatgataactaataacatagaacatttcaggaggacccttgg

Figure 2

## C. ELEGANS STRAIN

**BACTERIAL STRAIN**  
(only important phenotypes)

**VECTOR:**  
(only important features)




Figure 3.




Figure 4

# General description of the C.elegans T7 RNA polymerase expression vector with 4 examples




Figure 5





Figure 6



enhanced vector for RNAi, producing  
 sup35 dsRNA and dsRNA of the library, gene of interest  
 or PCR product

Figure 7




Figure 8




Figure 9








Figure 11




Figure 12




Figure 13




Figure 14

## pAS2\* with Forward and Reverse T7/T3/Sp6

T7/T3/Sp6 promotor (forward)

T7/T3/Sp6 promotor (reverse)




Figure 15

## pGAD424 with Forward and Reverse T7/T3/Sp6




Figure 16




Figure 17







Figure 18


let-858 promoter



a)

unc22-exon22

let-858 promoter



b)

Figure 19